

TWISTED STABILITY AND FOURIER-MUKAI TRANSFORM

KŌTA YOSHIOKA

0. INTRODUCTION

Let X be an abelian surface or a K3 surface over $\mathbb{C}$. Mukai introduced a lattice structure $\langle \ , \ \rangle$ on $H^{ev}(X, \mathbb{Z}) := \oplus H^{2i}(X, \mathbb{Z})$ by

$$(0.1) \quad \begin{aligned} \langle x, y \rangle &:= - \int_X x^\vee \wedge y \\ &= \int_X (x_1 \wedge y_1 - x_0 \wedge y_2 - x_2 \wedge y_0), \end{aligned}$$

where $x_i \in H^{2i}(X, \mathbb{Z})$ (resp. $y_i \in H^{2i}(X, \mathbb{Z})$) is the $2i$ -th component of x (resp. y) and $x^\vee = x_0 - x_1 + x_2$. It is now called Mukai lattice. For a coherent sheaf E on X , we can attach an element of $H^{ev}(X, \mathbb{Z})$ called Mukai vector $v(E) := \text{ch}(E)\sqrt{\text{td}_X}$, where $\text{ch}(E)$ is the Chern character of E and td_X is the Todd class of X . For a Mukai vector $v \in H^{ev}(X, \mathbb{Z})$ and an ample divisor H , let $M_H(v)$ be the moduli space of stable sheaves E of Mukai vector $v(E) = v$ and $\overline{M}_H(v)$ the moduli space of semi-stable sheaves. An ample divisor H is general with respect to v , if the following condition holds:

- (†) for every μ -semi-stable sheaf E of $v(E) = v$, if $F \subset E$ satisfies $(c_1(F), H)/\text{rk } F = (c_1(E), H)/\text{rk } E$, then $c_1(F)/\text{rk } F = c_1(E)/\text{rk } E$.

The preservation of stability by Fourier-Mukai transform on X was investigated by many people (e.g. [BBH2], [B-M], [Mu5], [Y8]). In [Y8], we introduced twisted degree of coherent sheaf E by $\deg_G(E) = \deg(E \otimes G^\vee)$, where G is a vector bundle on X . Then we showed that Fourier-Mukai transform preserves Gieseker semi-stability, if twisted degree is 0 and the polarization H is general. In this paper, we shall generalize our results to the case where H is not general. In this case, Fourier-Mukai transform does not preserve Gieseker semi-stability. This fact is closely related to the following fact: If H is not general, then Gieseker semi-stability is not preserved by the twisting $E \mapsto E \otimes L$, where L is a line bundle. Thus Gieseker semi-stability depends on the choice of L . In order to understand this phenomenon, Matsuki and Wentworth [M-W] (also by Ellingsrud and Göttsche [E-G] and Friedman and Qin [F-Q]) introduced L -twisted semi-stability, where L is a $\mathbb{Q}$ -line bundle. Hence we shall propose a formulation for our problem by using twisted semi-stability. In section 2, we shall show that Fourier-Mukai transform preserves suitable twisted semi-stability, if X is an abelian surface (Theorem 2.3).

In [Y8], we showed that $M_H(v)$ is deformation equivalent to a moduli space of torsion free sheaves of rank 1, if v is primitive and the polarization H is general. In section 3, we shall give another proof of this result by using results proved in section 2. Moreover we shall show the following.

Theorem 0.1. *Let X be an abelian surface or a K3 surface. Let $v \in H^{ev}(X, \mathbb{Z})$ be a Mukai vector of $\text{rk } v > 0$. Then $\overline{M}_H(v)$ is a normal variety, if $\langle v^2 \rangle > 0$ and H is general with respect to v .*

In section 4, we shall consider Fourier-Mukai transform on an Enriques surface associated to (-1) -reflection. In particular, we shall show a similar result to Theorem 2.3 (Proposition 4.3). As an application, we shall compute Hodge polynomials of some moduli spaces (Theorem 4.6). We also discuss a relation to Montonen-Olive duality in Physics (cf. [V-W]).

We are also interested in Fourier-Mukai transform on elliptic surfaces. Let $\pi : X \rightarrow C$ be an elliptic surface with a 0-section. Then a compactification of the relative Jacobian is isomorphic to X and there is a universal family $\mathcal{P}$ on $X \times_C X$. We regard $\mathcal{P}$ as a sheaf on $X \times X$ and consider Fourier-Mukai transform defined by $\mathcal{P}$. Assume that every fiber is irreducible. Then the preservation of stability was investigated in [Br1], [H-M], [J-M], [Y8]. In particular, semi-stable sheaf of relative degree 0 maps to a semi-stable sheaf of pure dimension 1. In [Br1], Bridgeland also treated Fourier-Mukai transform induced by a relative moduli space of stable sheaves on fibers. In order to generalize our result [Y8, Thm. 3.15] to this situation, we need to consider twisted semi-stability for purely 1-dimensional sheaves. In section 5, we introduce twisted stability for purely 1-dimensional sheaves and give some properties which are similar to results in [Y1]. Then we can show in Theorem 6.12 that a twisted semi-stable sheaf of twisted relative degree 0 maps to a twisted

semi-stable sheaf of pure dimension 1. As an application, we can compute Hodge numbers of some moduli spaces.

Theorem 0.2. *Let $\pi : X \rightarrow C$ be an elliptic surface with a section σ . Assume that every fiber is irreducible. f denotes a fiber of π . Let $M_{\sigma+kf}(r, c_1, \chi)$ be the moduli space of stable sheaves E of $(\text{rk}(E), c_1(E), \chi(E)) = (r, c_1, \chi)$ with respect to $\sigma + kf$. Then*

$$(0.2) \quad h^{p,q}(M_{\sigma+kf}(r, c_1, \chi)) = h^{p,q}(\text{Pic}^0(X) \times \text{Hilb}_X^n),$$

if $(r, (c_1, f)) = 1$ and $k \gg 0$, where $2n + h^1(\mathcal{O}_X) = \dim M_{\sigma+kf}(r, c_1, \chi)$.

Indeed we introduced twisted semi-stability for purely 1-dimensional sheaves to prove this theorem.

We also construct moduli spaces of twisted semi-stable sheaves by using Ellingsrud and Göttsche's method: They used moduli spaces of parabolic semi-stable torsion free sheaves constructed by Yokogawa [Yk]. Since his construction only works for parabolic semi-stable torsion free sheaves on smooth projective scheme, we need to construct moduli spaces of parabolic semi-stable sheaves of pure dimension 1, which will be done in appendix (Theorem 7.6).

1. PRELIMINARIES

Notation.

Throughout this note, we use the following notations. Let X be a smooth surface. For a scheme S , we denote the projection $S \times X \rightarrow S$ by p_S .

Let $\mathbb{Q}[[1/x, 1/y]]$ be the formal power series ring of two variable $1/x, 1/y$ and R the localization of $\mathbb{Q}[[1/x, 1/y]]$ by $1/(xy)$.

1.1. Virtual Hodge polynomial. For a variety Y over $\mathbb{C}$, cohomology with compact support $H_c^*(Y, \mathbb{Q})$ has a natural mixed Hodge structure. Let $e^{p,q}(Y) := \sum_k (-1)^k h^{p,q}(H_c^k(Y))$ be the virtual Hodge number and $e(Y) := \sum_{p,q} e^{p,q}(Y) x^p y^q$ the virtual Hodge polynomial of Y . For more details, see [Ch, 0.1]. By the properties of virtual Hodge polynomials, we can extend the definition of $e(Y)$ to constructible sets.

Let S be a bounded set of coherent sheaves on X with Hilbert polynomial $h(x)$. Under some conditions, we shall define the virtual Hodge polynomial of S as an element of R . So it is not a polynomial in general. Let H be an ample divisor on X . Since S is bounded, there is an integer m such that for every element E ,

- (i) $E(mH)$ is generated by global sections,
- (ii) $H^i(X, E(mH)) = 0$ for $i > 0$.

We set $N := h(m)$. We shall consider the quot scheme $Q := \text{Quot}_{\mathcal{O}_X(-mH)^{\oplus N}/X/\mathbb{C}}^{h(x)}$. Let $\mathcal{O}_{Q \times X}(-mH)^{\oplus N} \rightarrow Q$ be the universal quotient. We assume that

$$Q_S := \{q \in Q \mid \mathcal{Q}_q \in S, H^0(X, \mathcal{O}_X^{\oplus N}) \rightarrow H^0(X, \mathcal{Q}_q(mH)) \text{ is isomorphic}\}$$

is a constructible set. We shall show that this condition does not depend on the choice of Q . Let T be a scheme and $\mathcal{E}$ a coherent sheaf on $T \times X$ such that $\mathcal{E}$ is flat over T . Let T^0 be the open subscheme of T consisting of point $t \in T_S$ such that

- (i) $\mathcal{E}_t(mH)$ is generated by global sections,
- (ii) $H^i(X, \mathcal{E}_t(mH)) = 0$ for $i > 0$.

By base change theorem, $p_{T^0*}(\mathcal{E}(mH))$ is a locally free sheaf on T^0 . Let $\tau : \mathcal{P} \rightarrow T^0$ be the associated principal $GL(N)$ bundle over T^0 . Then there is a surjective homomorphism $\mathcal{O}_{\mathcal{P} \times X}(-mH)^{\oplus N} \rightarrow (\tau \times id_X)^* \mathcal{E}$. Hence we get a morphism $\eta : \mathcal{P} \rightarrow Q$. We set $T_S := \{t \in T^0 \mid \mathcal{E}_t \in S\}$. Then we see that $T_S = \tau(\eta^{-1}(Q_S))$. Thus T_S is constructible, if Q_S is constructible. In particular, constructibility does not depend on the choice of Q .

Definition 1.1. [Y6, Defn. 1.1]

- (i) S is constructible, if Q_S is constructible.
- (ii) For a constructible set S , we define the virtual Hodge polynomial by

$$e(S) := \frac{e(Q_S)}{e(GL(N))} \in R.$$

It is easy to see that this definition does not depend on the choice of Q . Let $\mathcal{M}_S$ be the associated substack of the stack of coherent sheaves. We define the virtual Hodge polynomial $e(\mathcal{M}_S)$ by $e(S)$.

1.2. Fourier-Mukai transform. Let $K(X)$ be the Grothendieck group of X . For $x \in K(X)$, we set

$$(1.1) \quad \gamma(x) := (\mathrm{rk} x, c_1(x), \chi(x)) \in \mathbb{Z} \oplus \mathrm{NS}(X) \oplus \mathbb{Z}.$$

Then $\gamma : K(X) \rightarrow \mathbb{Z} \oplus \mathrm{NS}(X) \oplus \mathbb{Z}$ is a surjective homomorphism and $\ker \gamma$ is generated by $\mathcal{O}_X(D) - \mathcal{O}_X$ and $\mathbb{C}_P - \mathbb{C}_Q$, where $D \in \mathrm{Pic}^0(X)$ and $P, Q \in X$.

For $\mathcal{E} \in \mathbf{D}(X_1 \times X_2)$, we define $\mathcal{F}_{\mathcal{E}} : \mathbf{D}(X_1) \rightarrow \mathbf{D}(X_2)$ by

$$(1.2) \quad \mathcal{F}_{\mathcal{E}}(x) := \mathbf{R}p_{2*}(\mathcal{E} \otimes p_1^*(x)), x \in \mathbf{D}(X_1).$$

and $\widehat{\mathcal{F}}_{\mathcal{E}} : \mathbf{D}(X_2) \rightarrow \mathbf{D}(X_1)$ by

$$(1.3) \quad \widehat{\mathcal{F}}_{\mathcal{E}}(y) := \mathbf{R} \mathrm{Hom}_{p_1}(\mathcal{E}, p_2^*(y)), y \in \mathbf{D}(X_2).$$

We denote the i -th cohomology $H^i(\mathcal{F}_{\mathcal{E}}(x))$ by $\mathcal{F}_{\mathcal{E}}^i(x)$.

$\mathcal{F}_{\mathcal{E}}$ also induces isomorphisms $\mathcal{F}_{\mathcal{E}} : K(X_1) \rightarrow K(X_2)$, $\mathcal{F}_{\mathcal{E}} : \mathbb{Z} \oplus \mathrm{NS}(X_1) \oplus \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathrm{NS}(X_2) \oplus \mathbb{Z}$ and we have a commutative diagram:

$$(1.4) \quad \begin{array}{ccc} \mathbf{D}(X_1) & \xrightarrow{\mathcal{F}_{\mathcal{E}}} & \mathbf{D}(X_2) \\ \downarrow & & \downarrow \\ K(X_1) & \xrightarrow{\mathcal{F}_{\mathcal{E}}} & K(X_2) \\ \gamma \downarrow & & \downarrow \gamma \\ \mathbb{Z} \oplus \mathrm{NS}(X_1) \oplus \mathbb{Z} & \xrightarrow{\mathcal{F}_{\mathcal{E}}} & \mathbb{Z} \oplus \mathrm{NS}(X_2) \oplus \mathbb{Z} \end{array}$$

For a divisor D , let $T_D : \mathbf{D}(X) \rightarrow \mathbf{D}(X)$ be an equivalence of derived categories defined by $T_D(x) = x \otimes \mathcal{O}_X(D)$. This is nothing but a Fourier-Mukai transform defined by $\mathcal{O}_{\Delta}(D)$, where $\Delta \subset X \times X$ is the diagonal.

Lemma 1.1. *Let $\mathcal{F}_{\mathcal{E}_1} : \mathbf{D}(X_1) \rightarrow \mathbf{D}(X_2)$ and $\mathcal{F}_{\mathcal{E}_2} : \mathbf{D}(X_2) \rightarrow \mathbf{D}(X_3)$ be Fourier-Mukai transforms defined by $\mathcal{E}_1 \in \mathbf{D}(X_1 \times X_2)$ and $\mathcal{E}_2 \in \mathbf{D}(X_2 \times X_3)$. Then we have $\mathcal{F}_{\mathcal{E}_2} \circ \mathcal{F}_{\mathcal{E}_1} = \mathcal{F}_{\mathcal{G}}$, where $\mathcal{G} := \mathcal{F}_{\mathcal{E}_2}(\mathcal{E}_1)$.*

1.3. Twisted stability for torsion free sheaves. Let H be an ample divisor on X . For $G \in K(X) \otimes \mathbb{Q}$ of $\mathrm{rk} G > 0$, we define G -twisted rank, degree, and Euler characteristic of $x \in K(X) \otimes \mathbb{Q}$ by

$$(1.5) \quad \begin{aligned} \mathrm{rk}_G(x) &:= \mathrm{rk}(G^{\vee} \otimes x) \\ \mathrm{deg}_G(x) &:= (c_1(G^{\vee} \otimes x), H) \\ \chi_G(x) &:= \chi(G^{\vee} \otimes x). \end{aligned}$$

For $t \in \mathbb{Q}_{>0}$, we get

$$(1.6) \quad \frac{\mathrm{deg}_G(x)}{\mathrm{rk}_G(x)} = \frac{\mathrm{deg}_{tG}(x)}{\mathrm{rk}_{tG}(x)}, \quad \frac{\chi_G(x)}{\mathrm{rk}_G(x)} = \frac{\chi_{tG}(x)}{\mathrm{rk}_{tG}(x)}.$$

We shall define G -twisted stability.

Definition 1.2. Let E be a torsion free sheaf on X . E is G -twisted semi-stable (resp. stable) with respect to H , if

$$(1.7) \quad \frac{\chi_G(F(nH))}{\mathrm{rk}_G(F)} \leq \frac{\chi_G(E(nH))}{\mathrm{rk}_G(E)}, n \gg 0$$

for $0 \subsetneq F \subsetneq E$ (resp. the inequality is strict).

For a $\mathbb{Q}$ -divisor α , we define α -twisted stability as $\mathcal{O}_X(\alpha)$ -twisted stability. This is nothing but the twisted stability introduced by Matsuki and Wentworth [M-W]. It is easy to see that G -twisted stability is determined by $\alpha = \det(G)/\mathrm{rk} G$. Hence G -twisted stability is the same as the Matsuki-Wentworth stability.

Definition 1.3. Let $\mathcal{M}_H^G(\gamma)^{ss}$ be the moduli stack of G -twisted semi-stable sheaves E of $\gamma(E) = \gamma$ and $\mathcal{M}_H^G(\gamma)^s$ the open substack consisting of G -twisted stable sheaves. For usual stability, i.e., $G = \mathcal{O}_X$, we denote $\mathcal{M}_H^{\mathcal{O}_X}(\gamma)^{ss}$ by $\mathcal{M}_H(\gamma)^{ss}$.

Remark 1.1. Let $c_1(G)/\mathrm{rk} G = aH + \beta$, $a \in \mathbb{Q}, \beta \in H^{\perp}$ be the orthogonal decomposition. Then the twisted semi-stability only depends on β , i.e., $\mathcal{M}_H^G(\gamma)^{ss} = \mathcal{M}_H^{\beta}(\gamma)^{ss}$.

Theorem 1.2. [M-W] (also see [E-G])

- (i) There is a coarse moduli scheme $\overline{\mathcal{M}}_H^G(\gamma)$ of S -equivalence classes of G -twisted semi-stable sheaves.
- (ii) $\overline{\mathcal{M}}_H^G(\gamma)$ is projective.

- (iii) For different G, G' , the relation between $\overline{M}_H^G(\gamma)$ and $\overline{M}_H^{G'}(\gamma)$ is described as Mumford-Thaddeus type flips:

$$(1.8) \quad \begin{array}{ccccccc} \overline{M}_H^{G_1}(\gamma) & & \overline{M}_H^{G_2}(\gamma) & & \overline{M}_H^{G_n}(\gamma) \\ & \searrow & \swarrow & \searrow & \swarrow & \dots & \swarrow \\ & \overline{M}_H^{G_{1,2}}(\gamma) & & \overline{M}_H^{G_{2,3}}(\gamma) & & & \end{array}$$

where $G = G_1, G' = G_n$.

Definition 1.4. $M_H^G(\gamma)$ is the open subscheme of $\overline{M}_H^G(\gamma)$ consisting of G -twisted stable sheaves and $M_H(\gamma)^{\mu-s}$ the open subscheme consisting of μ -stable sheaves. Usually we denote $\overline{M}_H^{\mathcal{O}_X}(\gamma)$ by $\overline{M}_H(\gamma)$ and $M_H^{\mathcal{O}_X}(\gamma)$ by $M_H(\gamma)$.

Since μ -stability does not depend on G , $M_H(\gamma)^{\mu-s}$ is a subscheme of $\overline{M}_H^G(\gamma)$ for all G .

Definition 1.5. For a pair (H, G) of an ample divisor H and an element $G \in K(X) \otimes \mathbb{Q}$, (H, G) is general with respect to v , if the following condition holds for every $E \in \mathcal{M}_H^G(v)^{ss}$:

For $0 \subsetneq F \subsetneq E$,

$$(1.9) \quad \frac{\chi_G(F(nH))}{\text{rk}_G(F)} = \frac{\chi_G(E(nH))}{\text{rk}_G(E)}, n \gg 0$$

implies that $v(F)/\text{rk } F = v/\text{rk } v$.

The following is easy (cf. [M-W]).

Lemma 1.3. For an ample divisor H and a Mukai vector v , there is a general (H, G) .

2. FOURIER-MUKAI TRANSFORM ON ABELIAN AND K3 SURFACES

Let X be a K3 surface or an abelian surface. Let E be a coherent sheaf. Let

$$(2.1) \quad \begin{aligned} v(E) &:= \text{ch}(E)\sqrt{\text{td}_X} \\ &= \text{rk}(E) + c_1(E) + (\chi(E) - \epsilon \text{rk}(E))\omega_X \in H^{ev}(X, \mathbb{Z}) \end{aligned}$$

be Mukai vector of E , where $\epsilon = 0, 1$ according as X is an abelian surface or a K3 surface and ω_X is the fundamental class of X . For these surfaces, it is common to use Mukai vector of E instead of using $\gamma(E)$. Hence we use Mukai vector in this section. For a Mukai vector v , we define $\mathcal{M}(v)$, $\mathcal{M}_H^G(v)$, $\overline{M}_H^G(v)$, $\dots$ as in section 1. By using twisted stability introduced by Matsuki and Wentworth [M-W], we shall generalize [Y8, sect. 8.2]. In order to state our theorem (Theorem 2.3), we prepare some notations.

Let $v_1 := r_1 + c_1 + a_1\omega_X, r_1 > 0, c_1 \in \text{NS}(X)$ be a primitive isotropic Mukai vector on X . We take a general ample divisor H with respect to v_1 . We set $Y := M_H(v_1)$. Assume that there is a universal family $\mathcal{E}$ on $X \times Y$. If X is an abelian surface, then Y consists of μ -stable vector bundles. By the proof of [Y7, Lem. 2.1], the following lemma holds.

Lemma 2.1. Assume that H is general with respect to v_1 .

- (i) If Y contains a non-locally free sheaf, then there is an exceptional vector bundle E_0 and $v_1 = \text{rk}(E_0)v(E_0) - \omega_X$. Moreover $Y \cong X$ and a universal family is given by

$$(2.2) \quad \mathcal{E} := \ker(E_0 \boxtimes E_0^\vee \rightarrow \mathcal{O}_\Delta).$$

- (ii) If Y consists of locally free sheaves, then they are μ -stable.

We set $w_1 := v(\mathcal{E}_{\{x\} \times Y}) = r_1 + \widehat{c}_1 + \widehat{a}_1\omega_Y, x \in X$. We consider a functor $\mathcal{H}_\mathcal{E} : \mathbf{D}(X) \rightarrow \mathbf{D}(Y)_{op}$ defined by

$$(2.3) \quad \mathcal{H}_\mathcal{E}(x) := \mathbf{R}\text{Hom}_{p_Y}(p_X^*(x), \mathcal{E}), x \in \mathbf{D}(X),$$

where $p_X : X \times Y \rightarrow X$ (resp. $p_Y : X \times Y \rightarrow Y$) be the projection. Then Bridgeland [Br2] showed that $\mathcal{H}_\mathcal{E}$ gives an equivalence of categories and the inverse is given by

$$(2.4) \quad \widehat{\mathcal{H}}_\mathcal{E}(y) := \mathbf{R}\text{Hom}_{p_X}(p_Y^*(y), \mathcal{E}), y \in \mathbf{D}(Y)_{op}.$$

$\mathcal{H}_\mathcal{E}$ induces an isometry $H^{ev}(X, \mathbb{Z}) \rightarrow H^{ev}(Y, \mathbb{Z})$. We also denote it by $\mathcal{H}_\mathcal{E}$.

We have an isomorphism $\text{NS}(X) \otimes \mathbb{Q} \rightarrow v_1^\perp \cap \omega_X^\perp$ by sending $D \in \text{NS}(X) \otimes \mathbb{Q}$ to $D + \frac{1}{r}(D, c_1)\omega_X \in v_1^\perp \cap \omega_X^\perp$. Since $\mathcal{H}_\mathcal{E}$ is an isometry of Mukai lattice, we get an isomorphism $v_1^\perp \cap \omega_X^\perp \rightarrow w_1^\perp \cap \omega_Y^\perp$. Thus we have an isomorphism $\delta : \text{NS}(X) \otimes \mathbb{Q} \rightarrow \text{NS}(Y) \otimes \mathbb{Q}$ given by

$$(2.5) \quad \delta(c_1(L)) = c_1(p_{Y*}(\text{ch } \mathcal{E} \sqrt{\text{td}_X} p_X^*(c_1(L) + \frac{1}{r}(c_1(L), c_1)\omega_X)^\vee)).$$

For a $\mathbb{Q}$ -line bundle $L \in \text{Pic}(X) \otimes \mathbb{Q}$, we choose a $\mathbb{Q}$ -line bundle $\widehat{L}$ on Y such that $\delta(c_1(L)) = c_1(\widehat{L})$. By a result of Li [Li] (or [BBH2]) and [Y8, Lem. 7.1], $\widehat{H}$ is ample, if Y consists of μ -stable vector bundles. By [Y7, Lem. 2.1], Y consists of μ -stable vector bundles unless $\mathcal{E}$ is given by (2.2). In this case, a direct computation (or [Li]) shows that $\widehat{L}$ is ample.

We consider the following two conditions.

- (#1) $\widehat{H}$ is general with respect to w_1 .
- (#2) $\mathcal{E}_{|\{x\} \times Y}$ is stable with respect to $\widehat{H}$.

Remark 2.1. If X is abelian or Y consists of non-locally free sheaves, then the assumption (#1, 2) holds for all general H . For another example, see [BBH1].

Problem. [Y8] Is $\mathcal{E}_{|\{x\} \times Y}$ always stable with respect to $\widehat{H}$?

For a coherent sheaf E on X (resp. F on Y), we set $\deg(E) := (c_1(E), H)$ (resp. $\deg(F) := (c_1(F), \widehat{H})$). We consider twisted degree $\deg_{G_1}(E)$ and $\deg_{G_2}(F)$, where $G_1 := \mathcal{E}_{|X \times \{y\}}$ and $G_2 := \mathcal{E}_{|\{x\} \times Y}$. Then

Lemma 2.2. [Y8, Lem. 8.3] $\deg_{G_1}(v) = \deg_{G_2}(\mathcal{H}_{\mathcal{E}}(v))$.

Every Mukai vector v is uniquely written as

$$(2.6) \quad v = lv_1 - a\omega_X + d(H + \frac{1}{r}(H, c_1)\omega_X) + (D + \frac{1}{r}(D, c_1)\omega_X),$$

where $l, a, d \in \mathbb{Q}$, and $D \in \text{NS}(X) \otimes \mathbb{Q} \cap H^\perp$.

It is easy to see that $l = -\langle v, \omega_X \rangle / \text{rk } v_1$, $a = \langle v, v_1 \rangle / \text{rk } v_1$ and $d = \deg_{G_1}(v) / r(H^2)$.

Definition 2.1. For a Mukai vector v , we set $l(v) := -\langle v, \omega_X \rangle / \text{rk } v_1$, $a(v) := \langle v, v_1 \rangle / \text{rk } v_1$.

Since $\mathcal{H}_{\mathcal{E}}(v_1) = \omega_Y$ and $\widehat{\mathcal{H}}_{\mathcal{E}}(w_1) = \omega_X$, we get

$$(2.7) \quad \mathcal{H}_{\mathcal{E}}(lv_1 - a\omega_X + (dH + D + \frac{1}{r}(dH + D, c_1)\omega_X)) = l\omega_Y - aw_1 + (d\widehat{H} + \widehat{D} + \frac{1}{r}(d\widehat{H} + \widehat{D}, \widehat{c}_1)\omega_Y)$$

where $\widehat{D} \in \text{NS}(X) \otimes \mathbb{Q} \cap H^\perp$. We can now state our theorem.

Theorem 2.3. We assume the condition (#1, 2) holds. Assume that $\deg_{G_1}(v) = 0$ and $l(v), a(v) > 0$. Let ε be an element of $K(X) \otimes \mathbb{Q}$ such that $v(\varepsilon) \in v_1^\perp \cap \omega_X^\perp$, $|\langle v(\varepsilon)^2 \rangle| \ll 1$ and $(H, c_1(\varepsilon)) = 0$. Then

$$(2.8) \quad \mathcal{M}_H^{G_1+\varepsilon}(v)^{ss} \rightarrow \mathcal{M}_{\widehat{H}}^{G_2+\widehat{\varepsilon}}(-\mathcal{H}_{\mathcal{E}}(v))^{ss}.$$

In particular, if $c_1(G_1) \in \mathbb{Q}H$, then $c_1(G_2) \in \mathbb{Q}\widehat{H}$ and we have an isomorphism $\mathcal{M}_H(v)^{ss} \rightarrow \mathcal{M}_{\widehat{H}}(-\mathcal{H}_{\mathcal{E}}(v))^{ss}$.

The proof of Theorem 2.3 is almost the same as that in [Y8, Thm. 8.2]. Before proving Theorem 2.3, we prepare two lemmas.

Lemma 2.4. Assume that $a(v) > 0$. Then $\text{Hom}(\mathcal{E}_{|X \times \{y\}}, E) = 0$ for all $y \in Y$ and $E \in \mathcal{M}_H^{G_1}(v)^{ss}$.

Proof. Since H is general with respect to v_1 , $\mathcal{E}_{|X \times \{y\}}$ is G_1 -twisted stable. Since E is G_1 -twisted semi-stable, it is sufficient to show that $-a(\mathcal{E}_{|X \times \{y\}}) / l(\mathcal{E}_{|X \times \{y\}}) > -a(v) / l(v)$. Since $v(\mathcal{E}_{|X \times \{y\}}) = v_1$, we get

$$(2.9) \quad \frac{-a(\mathcal{E}_{|X \times \{y\}})}{l(\mathcal{E}_{|X \times \{y\}})} - \frac{-a(v)}{l(v)} = \frac{a(v)}{l(v)} > 0.$$

□

Lemma 2.5. For a μ -semi-stable sheaf E of $v(E) = v$, there is a finite subset $S \subset Y$ such that

$$(2.10) \quad \text{Hom}(E, \mathcal{E}_{|X \times \{y\}}) = 0$$

for all $y \in Y \setminus S$.

Proof. Considering Jordan-Hölder filtration of E with respect to μ -stability, we may assume that E is μ -stable. If $\mathcal{E}_{|X \times \{y\}}$ is locally free, then by Lemma 2.1, $\mathcal{E}_{|X \times \{y\}}$ is μ -stable, and hence $E^{\vee\vee} \cong \mathcal{E}_{|X \times \{y\}}$. Therefore y is uniquely determined by E . Next we assume that $\mathcal{E}_{|X \times \{y\}}$ is not locally free. Under the notation of (2.2), if $E^{\vee\vee} \neq E_0$, then clearly $\text{Hom}(E, E_0) = 0$. Hence $\text{Hom}(E, \mathcal{E}_{|X \times \{y\}}) = 0$ for all $y \in Y$. If $E^{\vee\vee} = E_0$, then $\text{Hom}(E, \mathcal{E}_{|X \times \{y\}}) = 0$ for $y \in Y \setminus \text{Supp}(E^{\vee\vee}/E)$. □

Proof of Theorem 2.3. We shall first treat the case where $\varepsilon = 0$. By the symmetry of the condition, it is sufficient to show that WIT_1 holds for $E \in M_H(v)$ (i.e., $\mathcal{H}_\varepsilon^i(E) = 0, i \neq 1$) and $\mathcal{H}_\varepsilon^1(E)$ is G_2 -twisted semi-stable with respect to $\widehat{L}$. By Lemma 2.4 and 2.5, WIT_1 holds and $\mathcal{H}_\varepsilon^1(E)$ is torsion free. We shall show that E is G_2 -twisted semi-stable.

(I) $\mathcal{H}_\varepsilon^1(E)$ is μ -semi-stable: Assume that $\mathcal{H}_\varepsilon^1(E)$ is not μ -semi-stable. Let $0 \subset F_1 \subset F_2 \subset \cdots \subset F_s = \mathcal{H}_\varepsilon^1(E)$ be the Harder-Narasimhan filtration of $\mathcal{H}_\varepsilon^1(E)$ with respect to μ -semi-stability. We shall choose the integer k which satisfies $\deg_{G_2}(F_i/F_{i-1}) \geq 0, i \leq k$ and $\deg_{G_2}(F_i/F_{i-1}) < 0, i > k$. We claim that $\widehat{\mathcal{H}}_\varepsilon^0(F_k) = 0$ and $\widehat{\mathcal{H}}_\varepsilon^2(\mathcal{H}_\varepsilon^1(E)/F_k) = 0$. Indeed $\deg_{G_2}(F_i/F_{i-1}) \geq 0, i \leq k$ and the μ -semi-stability of F_i/F_{i-1} imply that $\widehat{\mathcal{H}}_\varepsilon^0(F_i/F_{i-1}), i \leq k$ is of dimension 0. Since $\widehat{\mathcal{H}}_\varepsilon^0(F_i/F_{i-1})$ is torsion free, $\widehat{\mathcal{H}}_\varepsilon^0(F_i/F_{i-1}) = 0, i \leq k$. Hence $\widehat{\mathcal{H}}_\varepsilon^0(F_k) = 0$. On the other hand, we also see that $\widehat{\mathcal{H}}_\varepsilon^2(F_i/F_{i-1}) = 0, i > k$. Hence we conclude that $\widehat{\mathcal{H}}_\varepsilon^2(\mathcal{H}_\varepsilon^1(E)/F_k) = 0$. So F_k and $\mathcal{H}_\varepsilon^1(E)/F_k$ satisfy WIT_1 and we get an exact sequence

$$(2.11) \quad 0 \rightarrow \widehat{\mathcal{H}}_\varepsilon^1(\mathcal{H}_\varepsilon^1(E)/F_k) \rightarrow E \rightarrow \widehat{\mathcal{H}}_\varepsilon^1(F_k) \rightarrow 0.$$

By (2.7), $\deg_{G_1}(\widehat{\mathcal{H}}_\varepsilon^1(F_k)) = -\deg_{G_2}(F_k) < 0$. This means that E is not μ -semi-stable with respect to L . Therefore $\mathcal{H}_\varepsilon^1(E)$ is μ -semi-stable with respect to L .

(II) $\mathcal{H}_\varepsilon^1(E)$ is G_2 -twisted semi-stable: Assume that $\mathcal{H}_\varepsilon^1(E)$ is not G_2 -twisted semi-stable. Then there is an exact sequence

$$(2.12) \quad 0 \rightarrow F_1 \rightarrow \mathcal{H}_\varepsilon^1(E) \rightarrow F_2 \rightarrow 0$$

such that (i) F_2 is G_2 -twisted stable and (ii) $-a(F_2)/l(F_2) < -a(\mathcal{H}_\varepsilon^1(E))/l(\mathcal{H}_\varepsilon^1(E)) = -l(v)/a(v)$, where $v(F_2) = l(F_2)w_1 - a(F_2)\omega_Y$. Since $-a(F_2)/l(F_2) < -l(v)/a(v) < 0$, Lemma 2.4 and 2.5 imply that $\widehat{\mathcal{H}}_\varepsilon^0(F_2) = \widehat{\mathcal{H}}_\varepsilon^2(F_2) = 0$. We also obtain that $\widehat{\mathcal{H}}_\varepsilon^0(F_1) = 0$. Hence we have an exact sequence

$$(2.13) \quad 0 \rightarrow \widehat{\mathcal{H}}_\varepsilon^1(F_2) \rightarrow E \rightarrow \widehat{\mathcal{H}}_\varepsilon^1(F_1) \rightarrow 0.$$

Since $\widehat{\mathcal{H}}_\varepsilon^1(\mathcal{H}_\varepsilon^1(E)) = E$, $\widehat{\mathcal{H}}_\varepsilon^2(F_1) = 0$. Thus WIT_1 also holds for F_1 . By (ii), we see that

$$(2.14) \quad \begin{aligned} \frac{-a(v)}{l(v)} - \frac{-a(\widehat{\mathcal{H}}_\varepsilon^1(F_2))}{l(\widehat{\mathcal{H}}_\varepsilon^1(F_2))} &= \frac{-a(v)}{l(v)} + \frac{l(F_2)}{a(F_2)} \\ &= \frac{-a(v)a(F_2) + l(v)l(F_2)}{l(v)a(F_2)} < 0. \end{aligned}$$

This means that E is not G_1 -twisted semi-stable. Therefore $\mathcal{H}_\varepsilon^1(E)$ is G_2 -twisted semi-stable.

We next treat general cases. Since $|\langle \varepsilon^2 \rangle| \ll 1$, we have an inclusion $\mathcal{M}_H^{G_1+\varepsilon}(v)^{ss} \subset \mathcal{M}_H^{G_1}(v)^{ss}$ and the complement consists of E which fits in an exact sequence:

$$(2.15) \quad 0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0$$

where E_1 is a G_1 -twisted semi-stable sheaf such that $v(E_1) = l_1 v_1 - a_1 \omega + \delta_1$, $\delta_1 \in v_1^\perp \cap \omega_X^\perp \cap H^\perp$, $a_1/l_1 = a(v)/l(v)$ and $-\langle v(E_1), v_1 + \epsilon \rangle/l_1 > -\langle v, v_1 + \epsilon \rangle/l(v)$. Then we see that $-\langle \delta_1, \epsilon \rangle/l_1 > -\langle \delta, \epsilon \rangle/l(v)$, where $\delta := v - (l(v)v_1 - a(v)\omega_X) \in v_1^\perp \cap \omega_X^\perp \cap H^\perp$. Applying $\mathcal{H}_\varepsilon^1$ to the exact sequence (2.15), we get an exact sequence

$$(2.16) \quad 0 \rightarrow \mathcal{H}_\varepsilon^1(E_2) \rightarrow \mathcal{H}_\varepsilon^1(E) \rightarrow \mathcal{H}_\varepsilon^1(E_1) \rightarrow 0.$$

Since $-\langle \widehat{\delta}_1, \widehat{\epsilon} \rangle/a_1 > -\langle \widehat{\delta}, \widehat{\epsilon} \rangle/a(v)$, we get that $-\langle v(\mathcal{H}_\varepsilon^1(E_1)), \widehat{\epsilon} \rangle/a_1 < -\langle v(\mathcal{H}_\varepsilon^1(E)), \widehat{\epsilon} \rangle/a(v)$. Therefore $\mathcal{H}_\varepsilon^1(E)$ is not $(G_2 + \widehat{\epsilon})$ -twisted semi-stable. $\square$

Proposition 2.6. Assume that $M_H(v)^{\mu-s}$ is an open dense subscheme of $\overline{M}_H(v)$. If $\deg_{G_1}(v) = 0$, then $\mathcal{H}_\varepsilon$ induces a birational map $\overline{M}_H(v) \cdots \rightarrow \overline{M}_H(\mathcal{H}_\varepsilon(v))$ which is described as Mumford-Thaddeus type flips:

$$(2.17) \quad \begin{array}{ccccccc} \overline{M}_H^{\alpha_1}(v) & & \overline{M}_H^{\alpha_2}(v) & & \overline{M}_H^{\alpha_n}(\mathcal{H}_\varepsilon(v)) \\ & \searrow & \swarrow & \searrow & \cdots & \swarrow \\ & \overline{M}_H^{\alpha_{1,2}}(v) & & \overline{M}_H^{\alpha_{2,3}}(v) & & \end{array}$$

where $\alpha_i, \alpha_{i,i+1} \in \text{NS}(X) \otimes \mathbb{Q}$ and $\alpha_1 = \alpha_n = 0$.

Proof. By Theorem 1.2, we have Mumford-Thaddeus type flips $\overline{M}_H(v) \cdots \rightarrow \overline{M}_H^{G_1}(v)$ and $\overline{M}_H^{G_2}(\mathcal{H}_\varepsilon(v)) \cdots \rightarrow \overline{M}_H^{G_1}(\mathcal{H}_\varepsilon(v))$. By Theorem 2.3, $\overline{M}_H^{G_1}(v) \cong \overline{M}_H^{G_2}(\mathcal{H}_\varepsilon(v))$. Therefore we get our claim. $\square$

Example 2.1. Let X be a K3 surface and H an ample divisor on X . Assume that $H^\perp = \mathbb{Z}D$ and $(D^2) = -2n$, $n > 2$. We set $v = 2 + (1 - 2n)\omega_X$. Then there is a non-trivial extension

$$(2.18) \quad 0 \rightarrow I_x(D) \rightarrow E \rightarrow \mathcal{O}_X(-D) \rightarrow 0$$

where $x \in X$. We can easily show that E is a stable sheaf of $v(E) = v$. We consider Fourier-Mukai transform defined by $\mathcal{E} = I_\Delta \otimes p_X^* \mathcal{O}_X(D)$. Since $\text{Ext}^2(E, \mathcal{E}_x) = \text{Hom}(\mathcal{E}_x, E)^\vee \neq 0$, E does not satisfy WIT_1 with respect to $\mathcal{H}_\mathcal{E}$. In this case, we get the following diagram

$$(2.19) \quad \begin{array}{ccc} \overline{M}_H(v) & \cdots \rightarrow & \overline{M}_H^D(v) \cong \overline{M}_H(\mathcal{H}_\mathcal{E}(v)) \\ & \searrow & \swarrow \\ & \overline{M}_H^{tD}(v) & \end{array}$$

where $t = 1/4n$.

Remark 2.2. Let (X, H) be a polarized K3 surface which has a divisor ℓ such that

$$(2.20) \quad (H^2) = 2, (\ell^2) = -12, (H, \ell) = 0$$

and $H^0(X, \mathcal{O}_X(\ell + 2H)) = 0$. Then $Y := M_H(2 + \ell - 3\omega_X)$ is isomorphic to X and there is a universal family $\mathcal{E}$ on $X \times Y$. In [B-M], Bruzzo and Maciocia showed that Fourier-Mukai transform $\mathcal{F}_\mathcal{E}$ gives an isomorphism

$$(2.21) \quad \mathcal{M}_H(1 + (1 - n)\omega_X)^{ss} \cong \mathcal{M}_{\widehat{H}}((1 + 2n) - n\widehat{\ell} + (1 - 3n)\omega_Y)^{ss}.$$

Moreover every element E of $\mathcal{M}_{\widehat{H}}((1 + 2n) - n\widehat{\ell} + (1 - 3n)\omega_Y)$ fits in a non-trivial extension

$$(2.22) \quad 0 \rightarrow E' \rightarrow E \rightarrow \mathcal{O}_Y \rightarrow 0$$

where $E' \in \mathcal{M}_{\widehat{H}}(n(2 - \widehat{\ell} - 3\omega_Y))^{ss}$. Then we can show that $E \mapsto E^\vee$ induces an isomorphism $\mathcal{M}_{\widehat{H}}((1 + 2n) - n\widehat{\ell} + (1 - 3n)\omega_Y)^{ss} \rightarrow \mathcal{M}_{\widehat{H}}^{\widehat{\ell}/2}((1 + 2n) + n\widehat{\ell} + (1 - 3n)\omega_Y)^{ss}$. Thus we get an isomorphism

$$(2.23) \quad \mathcal{M}_H(1 + (1 - n)\omega_X)^{ss} \cong \mathcal{M}_{\widehat{H}}^{\widehat{\ell}/2}((1 + 2n) + n\widehat{\ell} + (1 - 3n)\omega_Y)^{ss},$$

which is nothing but the isomorphism given by $\mathcal{H}_{\mathcal{E}^\vee}$.

3. IRREDUCIBILITY OF $\overline{M}_H(v)$

3.1. A special case of Theorem 2.3. We shall give an application of Theorem 2.3. Let X be an abelian surface or a K3 surface such that $\text{NS}(X) = \mathbb{Z}e \oplus \mathbb{Z}f$, $(e^2) = (f^2) = 0$ and $(e, f) = 1$.

Corollary 3.1. *Assume that (1) X is an abelian surface, Y is the dual abelian surface and $\mathcal{E}$ is the Poincaré line bundle on $X \times Y$, or (2) X is a K3 surface, $Y = X$ and $\mathcal{E}$ is the ideal sheaf of the diagonal $\Delta \subset X \times X$. Assume that $e + kf$ is an ample divisor. We set $D := e - kf$. Then $\mathcal{H}_\mathcal{E}$ induces an isomorphism of stacks*

$$(3.1) \quad \mathcal{M}_{e+kf}(r + cD - a\omega_X)^{ss} \rightarrow \mathcal{M}_{\widehat{e}+k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss},$$

where $r, a > 0$ and $c \geq 0$. Moreover, if k is a sufficiently large integer depending on r and $\langle (r + cD - a\omega_X)^2 \rangle$, then

$$(3.2) \quad \mathcal{M}_{e+nf}(r + cD - a\omega_X)^{ss} \cong \mathcal{M}_{\widehat{e}+n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$$

if $0 < n - k \ll 1$, where $n \in \mathbb{Q}$.

Remark 3.1. By Hodge index theorem, $(D^2) < 0$. Hence if $c > 0$, then $a > 0$, or $a = 0$ and $(D^2) = -2$.

Proof. By Theorem 2.3, we get the first claim. We next show the second claim. We note that $(D^2) = -2k \ll 0$. Hence $e + kf$ is a general polarization with respect to $r + cD - a\omega_X$ (cf. [Y1, Lem. 2.1, Rem. 2.1]). The same is true for $e + nf$, $n > k$. Hence $\mathcal{M}_{e+kf}(r + cD - a\omega_X)^{ss} \cong \mathcal{M}_{e+nf}(r + cD - a\omega_X)^{ss}$. In order to prove our claim, it suffices to show that $\mathcal{M}_{\widehat{e}+n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss} = \mathcal{M}_{\widehat{e}+k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$. We first show that $\mathcal{M}_{\widehat{e}+k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss} \subset \mathcal{M}_{\widehat{e}+n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$. Assume that there is an exact sequence

$$(3.3) \quad 0 \rightarrow F_1 \rightarrow \mathcal{H}_\mathcal{E}^1(E) \rightarrow F_2 \rightarrow 0$$

such that F_2 is semi-stable with respect to $\widehat{e} + n\widehat{f}$ and

$$(3.4) \quad \frac{(c_1(\mathcal{H}_\mathcal{E}^1(E)), \widehat{e} + n\widehat{f})}{\text{rk } \mathcal{H}_\mathcal{E}^1(E)} > \frac{(c_1(F_2), \widehat{e} + n\widehat{f})}{\text{rk } F_2}$$

or

$$(3.5) \quad \frac{(c_1(\mathcal{H}_\mathcal{E}^1(E)), \widehat{e} + n\widehat{f})}{\text{rk } \mathcal{H}_\mathcal{E}^1(E)} = \frac{(c_1(F_2), \widehat{e} + n\widehat{f})}{\text{rk } F_2}, \quad \frac{\chi(\mathcal{H}_\mathcal{E}^1(E))}{\text{rk } \mathcal{H}_\mathcal{E}^1(E)} > \frac{\chi(F_2)}{\text{rk } F_2}.$$

Since $0 < n - k \ll 1$, (i) or (ii) implies that F_1 and F_2 are μ -semi-stable of $(c_1(F_1), \widehat{e} + k\widehat{f}) = (c_1(F_2), \widehat{e} + k\widehat{f}) = 0$ with respect to $\widehat{e} + k\widehat{f}$. Hence $\text{Hom}(F_1, \mathcal{E}_x) = \text{Hom}(F_2, \mathcal{E}_x) = 0$ except finite number of points of X .

If (i) holds, then $\text{Ext}^2(F_2, \mathcal{E}_{|X \times \{y\}}) = \text{Hom}(\mathcal{E}_{|X \times \{y\}}, F_2)^\vee = 0$ for all $y \in Y$, because $\mathcal{E}_{|X \times \{y\}}$, $y \in Y$ is a stable sheaf of $c_1(\mathcal{E}_{|X \times \{y\}}) = 0$ with respect to $e + nf$ and $(c_1(F_2), \widehat{e} + n\widehat{f})/\text{rk } F_2 < (-c\widehat{D}, \widehat{e} + n\widehat{f})/\text{rk } \mathcal{H}_\mathcal{E}^1(E) = -c(n - k)/\text{rk } \mathcal{H}_\mathcal{E}^1(E) \leq 0$. Therefore F_1 and F_2 satisfies WIT_1 and we get an exact sequence

$$(3.6) \quad 0 \rightarrow \widehat{\mathcal{H}}_\mathcal{E}^1(F_2) \rightarrow E \rightarrow \widehat{\mathcal{H}}_\mathcal{E}^1(F_1) \rightarrow 0.$$

By Lemma 2.2,

$$(3.7) \quad (0 <) \frac{(c_1(E), e + nf)}{\text{rk } \mathcal{H}_\mathcal{E}^1(E)} < \frac{(c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2)), e + nf)}{\text{rk } F_2}.$$

Since E is semi-stable with respect to $e + kf$ and $(c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2)), e + kf) = 0$, $-\text{rk}(F_2)/\text{rk } \widehat{\mathcal{H}}_\mathcal{E}^1(F_2) \leq -\text{rk } \mathcal{H}_\mathcal{E}^1(E)/\text{rk } E$. Hence we see that

$$(3.8) \quad \frac{(c_1(E), e + nf)}{\text{rk } E} < \frac{(c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2)), e + nf)}{\text{rk } \widehat{\mathcal{H}}_\mathcal{E}^1(F_2)}.$$

This implies that E is not semi-stable with respect to $e + nf$. Therefore (i) does not occur. If (ii) holds, then $\frac{c_1(\mathcal{H}_\mathcal{E}^1(E))}{\text{rk } \mathcal{H}_\mathcal{E}^1(E)} = \frac{c_1(F_2)}{\text{rk } F_2}$. By the proof of Theorem 2.3, we get a contradiction. Thus $\mathcal{M}_{\widehat{e} + k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss} \subset \mathcal{M}_{\widehat{e} + n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$.

We next show that $\mathcal{M}_{\widehat{e} + n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss} \subset \mathcal{M}_{\widehat{e} + k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$. Assume that there is an element $F \in \mathcal{M}_{\widehat{e} + n\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss} \setminus \mathcal{M}_{\widehat{e} + k\widehat{f}}(a - c\widehat{D} - r\omega_Y)^{ss}$. Then we see that there is an exact sequence

$$(3.9) \quad 0 \rightarrow F_1 \rightarrow F \rightarrow F_2 \rightarrow 0$$

such that (i) $(c_1(F), \widehat{e} + n\widehat{f})/\text{rk } F \leq (c_1(F_2), \widehat{e} + n\widehat{f})/\text{rk } F_2$, (ii) $(c_1(F), \widehat{e} + k\widehat{f})/\text{rk } F = (c_1(F_2), \widehat{e} + k\widehat{f})/\text{rk } F_2$, (iii) $\chi(F)/\text{rk } F > \chi(F_2)/\text{rk } F_2$ and (iv) F_2 is semi-stable with respect to $\widehat{e} + k\widehat{f}$. We set $\epsilon = 0, 1$ according as X is an abelian surface or a K3 surface as in section 2. We note that

- (a) $\mathcal{E}_{|\{x\} \times Y}$, $x \in X$ is stable of $(c_1(\mathcal{E}_{|\{x\} \times Y}), \chi(\mathcal{E}_{|\{x\} \times Y})) = (0, \epsilon)$ with respect to $\widehat{e} + n\widehat{f}$,
- (b) F is stable of $(c_1(F), \widehat{e} + n\widehat{f}) \leq 0$ and $\chi(F)/\text{rk } F = \epsilon - a/r < \epsilon$ with respect to $\widehat{e} + n\widehat{f}$.

By (a) and (b), we get $\text{Ext}^2(F, \mathcal{E}_{|\{x\} \times Y}) = \text{Hom}(\mathcal{E}_{|\{x\} \times Y}, F)^\vee = 0$ for all $x \in X$. By (iii) and (iv), we see that $\text{Ext}^2(F_2, \mathcal{E}_{|\{x\} \times Y}) = 0$ for all $x \in X$. Since F_1 and F_2 are μ -semi-stable sheaves of degree 0 with respect to $\widehat{e} + k\widehat{f}$, $\text{Hom}(F_1, \mathcal{E}_{|\{x\} \times Y}) = \text{Hom}(F_2, \mathcal{E}_{|\{x\} \times Y}) = 0$ except finite number of points $x \in X$. Hence WIT_1 holds for F_1 , F_2 and F with respect to $\widehat{\mathcal{H}}_\mathcal{E}$ and we have an exact sequence

$$(3.10) \quad 0 \rightarrow \widehat{\mathcal{H}}_\mathcal{E}^1(F_2) \rightarrow \widehat{\mathcal{H}}_\mathcal{E}^1(F) \rightarrow \widehat{\mathcal{H}}_\mathcal{E}^1(F_1) \rightarrow 0.$$

In the same way, we see that $E := \widehat{\mathcal{H}}_\mathcal{E}^1(F)$ is μ -semi-stable with respect to $e + kf$. Since $e + kf$ is general, we get $c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2))/\text{rk } \widehat{\mathcal{H}}_\mathcal{E}^1(F_2) = c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F))/\text{rk } \widehat{\mathcal{H}}_\mathcal{E}^1(F)$. On the other hand, (i) implies that

$$(3.11) \quad \frac{(c_1(E), e + nf)}{\text{rk } F} \geq \frac{(c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2)), e + nf)}{\text{rk } F_2}.$$

By using (iii), we see that

$$(3.12) \quad \frac{(c_1(E), e + nf)}{\text{rk } E} > \frac{(c_1(\widehat{\mathcal{H}}_\mathcal{E}^1(F_2)), e + nf)}{\text{rk } \widehat{\mathcal{H}}_\mathcal{E}^1(F_2)},$$

which is a contradiction. Therefore our claim holds. $\square$

Remark 3.2. If $e + kf$ is general with respect to $r + cD - a\omega_X$ and $c > 0$, then $\text{Hom}(E, \mathcal{E}_x) = 0$ for all $x \in X$ and $E \in \mathcal{M}_{e + kf}(r + cD - a\omega_X)^{ss}$. Hence $\mathcal{H}_\mathcal{E}^1(E)$ is locally free.

Remark 3.3. In general, $\widehat{e} + k\widehat{f}$ is not a general polarization with respect to $a - c\widehat{D} - r\omega_Y$. Indeed, let E be a non-locally free μ -stable sheaf of $v(E) = r + cD - a\omega_X$ on X . Assume that $E^{\vee\vee}/E = \mathbb{C}_x$, $x \in X$ and $a > 1$. Then we get an exact sequence

$$(3.13) \quad 0 \rightarrow \mathcal{H}_\mathcal{E}^1(E^{\vee\vee}) \rightarrow \mathcal{H}_\mathcal{E}^1(E) \rightarrow \mathcal{H}_\mathcal{E}^2(\mathbb{C}_x) \rightarrow 0.$$

It is easy to see that $\mathcal{H}_\mathcal{E}^2(\mathbb{C}_x) \cong \mathcal{E}_x$. Hence $\widehat{e} + k\widehat{f}$ is not general with respect to $a - c\widehat{D} - r\omega_Y$, if $c > 0$.

3.2. Application to the deformation type of $M_H(v)$. Let X_1 be an abelian surface or a K3 surface.

Definition 3.1. Let v be a Mukai vector of $\text{rk } v > 0$. Then we can write it as $v = m(v)v_p$, where $m(v) \in \mathbb{Z}$ and v_p is a primitive Mukai vector of $\text{rk } v_p > 0$.

In [Y8], we showed that $M_H(v)$ is deformation equivalent to a moduli space of rank 1 torsion free sheaves, if v is primitive. Here we assume that $\text{rk } v > 0$ and H is general. We shall give a slightly different proof of this result, that is, we shall use O'Grady's arguments [O1, sect. 2]. One of the benefit of O'Grady's arguments is that we do not need to use algebraic space, which enable us to treat non-primitive Mukai vector cases. For a Mukai vector $v := l(r + c_1) + a\omega_{X_1} \in H^*(X_1, \mathbb{Z})$ such that $r > 0$, $l = \gcd(r, c_1)$ and $\gcd(l, a) = 1$, we set $b = -a + l\lambda$, $k = -(c_1^2)/2 + r\lambda$, $\lambda \gg 0$ so that $e + kf$ is ample. We consider X in the above notation. By [Y4, Prop. 1.1] or a modification of its proof, we see that $M_H(l(r + c_1) + a\omega_{X_1})$ is deformation equivalent to $M_{e+nf}(l(r + (e - kf)) - b\omega_X)$, where H is general with respect to v and $0 < n - k \ll 1$. By Corollary 3.1, we have an isomorphism

$$(3.14) \quad M_{e+nf}(l(r + (e - kf)) - b\omega_X) \rightarrow M_{\widehat{e}+n\widehat{f}}(b - l(\widehat{e} - k\widehat{f}) - l\omega_Y).$$

Since $(b, l) = 1$, $M_{\widehat{e}+n\widehat{f}}(b - l(\widehat{e} - k\widehat{f}) - l\omega_Y)$ is deformation equivalent to $M_{e+nf}(b + (e - k'f) - b'\omega_X)$, where $b' = lr + \lambda'$, $k' = l^2k + b\lambda'$, $\lambda' \gg 0$ and $0 < n - k' \ll 1$. Applying Corollary 3.1 again, we get an isomorphism

$$(3.15) \quad M_{e+nf}(b + (e - k'f) - b'\omega_X) \rightarrow M_{\widehat{e}+n\widehat{f}}(b' - (\widehat{e} - k'\widehat{f}) - b\omega_Y).$$

If λ' is sufficiently large, then $M_{\widehat{e}+n\widehat{f}}(b' - (\widehat{e} - k'\widehat{f}) - b\omega_Y)$ is deformation equivalent to $M_{\widehat{e}+n\widehat{f}}(b' - (\widehat{e} - k''\widehat{f}) - \omega_Y)$ and $e + k''f$ is ample, where $k'' = lr(1 - b) + l^2k + \lambda'$ and $0 < n - k \ll 1$. Since $k'' \gg 0$, Corollary 3.1 implies that $M_{e+nf}(1 + (e - k''f) - b''\omega_X)$ is isomorphic to $M_{\widehat{e}+n\widehat{f}}(b' - (\widehat{e} - k''\widehat{f}) - \omega_Y)$. Therefore $M_H(v)$ is deformation equivalent to the moduli space of rank 1 torsion free sheaves.

We shall next treat non-primitive Mukai vector.

Lemma 3.2. *Let v be a Mukai vector of $\text{rk } v > 0$. Let H be a general ample divisor with respect to v . We set*

$$(3.16) \quad \mathcal{M}_H(v)^{pss} := \{E \in \mathcal{M}_H(v)^{ss} \mid E \text{ is properly semi-stable}\}.$$

Then $\dim \mathcal{M}_H(v)^{pss} \leq \langle v^2 \rangle$. Moreover inequality is strict, unless $m(v) = 2$ and $\langle v^2 \rangle = 8$.

For the proof, see [Y7, Lem. 1.7].

Proposition 3.3. *Under the same assumptions, $\mathcal{M}_H(v)^{ss}$ is a locally complete intersection stack which contains $\mathcal{M}_H(v)^s$ as an open dense substack and the singular locus is at least of codimension 2. In particular $\mathcal{M}_H(v)^{ss}$ is normal.*

Proof. By deformation theory and Lemma 3.2, we see that $\mathcal{M}_H(v)^{ss}$ is a locally complete intersection. If $m(v) \neq 2$ or $\langle v^2 \rangle > 8$, then $\dim \mathcal{M}_H(v)^{pss} \leq \langle v^2 \rangle - 1$. Therefore the singular locus is at least of codimension 2. If $m(v) = 2$ and $\langle v^2 \rangle = 8$, then a general member of $\mathcal{M}_H(v)^{pss}$ fits in a non-trivial extension

$$(3.17) \quad 0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0$$

where $E_1, E_2 \in \mathcal{M}_H(v/2)^{ss}$ and $E_1 \neq E_2$. Then E is simple, which implies that $\mathcal{M}_H(v)^{ss}$ is smooth at E . Therefore the singular locus is at least of codimension 2. For the last claim, we use Serre's criterion. $\square$

Definition 3.2. Let Y_1, Y_2 be normal schemes. Then $Y_1 \sim Y_2$, if there is a proper and flat morphism $\mathcal{Y} \rightarrow T$ over a smooth connected curve T such that every fiber is normal and $Y_i = \mathcal{Y}_{t_i}$ for some $t_1, t_2 \in T$. Deformation equivalence is an equivalence relation generated by $\sim$.

By the following lemma, the number of irreducible components is an invariant of this equivalence relation.

Lemma 3.4. *Let T be a smooth curve and $Y \rightarrow T$ a flat and proper morphism. Assume that every fiber is normal. Then the number of irreducible components of Y_t , $t \in T$ is constant.*

Proof. Since Y_t is normal, every connected component is an integral scheme. Hence the number of irreducible components of Y_t is $h^0(Y_t, \mathcal{O}_{Y_t})$. By the upper-semicontinuity of $h^0(Y_t, \mathcal{O}_{Y_t})$, the number of irreducible components of Y_t is upper semi-continuous. On the other hand, by Zariski's connectivity theorem, the number of connected components of Y_t is lower semi-continuous. Therefore we get our lemma. $\square$

By the same proof, we can show the following.

Proposition 3.5. *$\overline{M}_H(v)$ is deformation equivalent to $\overline{M}_H(m(v)(1 - n\omega_X))$, where $n = \langle v_p^2 \rangle / 2$. In particular the number of irreducible components of $\overline{M}_H(v)$ is determined by $m(v)$.*

Proof. Let T be a smooth curve over $\mathbb{C}$ and $\varphi : (\mathcal{X}, \mathcal{L}) \rightarrow T$ be a family of polarized abelian or K3 surfaces. For a family of Mukai vectors $v \in R^*\varphi_*\mathbb{Z} = \cup_{t \in T} H^*(\mathcal{X}_t, \mathbb{Z})$, let $\psi : \overline{\mathfrak{M}}_{\mathcal{L}}(v) \rightarrow T$ be the relative moduli space of semi-stable sheaves on $\mathcal{X}_t$, $t \in T$ of Mukai vector v_t and $\mathfrak{M}_{\mathcal{L}}(v)$ the open subscheme consisting of stable sheaves. Since T is defined over a field of characteristic 0, $\overline{\mathfrak{M}}_{\mathcal{L}}(v)_t = \overline{M}_{\mathcal{L}_t}(v_t)$ for $t \in T$, where $\overline{M}_{\mathcal{L}_t}(v_t)$ is the moduli space of semi-stable sheaves on $\mathcal{X}_t$ (cf. [MFK, Thm. 1.1]). Since $\psi|_{\mathfrak{M}_{\mathcal{L}}(v)} : \mathfrak{M}_{\mathcal{L}}(v) \rightarrow T$ is smooth [Mu3], it is flat. Assume that $\mathcal{L}_t$ is general with respect to v_t for all $t \in T$. By Proposition 3.3, $\mathfrak{M}_{\mathcal{L}}(v)$ is a dense subscheme of $\overline{\mathfrak{M}}_{\mathcal{L}}(v)$. Since T is a smooth curve, ψ is also flat. Therefore $\psi : \overline{\mathfrak{M}}_{\mathcal{L}}(v) \rightarrow T$ is a proper and flat morphism. Then our claim follows from the same argument as in $m(v) = 1$ case. $\square$

3.3. Irreducibility of $\overline{M}_H(v)$. We shall show that $\overline{M}_H(v)$ is irreducible. We may assume that X has an elliptic fibration $\pi : X \rightarrow C$. We also assume that there is a section σ of π and $\text{NS}(X) = \mathbb{Z}\sigma \oplus \mathbb{Z}f$, where f is a fiber of π . By [Y8, Thm. 3.15] (cf. Theorem 6.12), we have an isomorphism $\overline{M}_{\sigma+kf}(r(1-n\omega_X)) \cong \overline{M}_{\sigma+kf}(w)$, where $w = r((\sigma + (n+\epsilon)f) + \omega_X)$. Hence it is sufficient to show that $\overline{M}_{\sigma+kf}(w)$ is irreducible. From now on, we assume that $r \geq 2$. By Proposition 3.3, we shall show that $M_{\sigma+kf}(w)$ is irreducible.

Definition 3.3. For a purely 1-dimensional sheaf L on X , $\text{Div}(L)$ is the divisor on X which is defined by the fitting ideal of L .

We set $\xi = r(\sigma + (n+\epsilon)f)$. Let Hilb_X^ξ be the Hilbert scheme of curves C on X such that $c_1(\mathcal{O}_X(C)) = \xi$. There is a natural map $j : \overline{M}_{\sigma+kf}(w) \rightarrow \text{Hilb}_X^\xi$ sending $L \in \overline{M}_{\sigma+kf}(w)$ to $\text{Div}(L)$. We want to estimate the dimension of closed subsets of $M_{\sigma+kf}(w)$:

$$(3.18) \quad \begin{aligned} N_1 &:= \{L \in M_{\sigma+kf}(w) \mid \text{Div}(L) \text{ is not irreducible}\}, \\ N_2 &:= \{L \in M_{\sigma+kf}(w) \mid \text{Div}(L) \text{ is not reduced}\}. \end{aligned}$$

Estimate of $\dim N_1$. We prepare some lemmas.

Lemma 3.6. *Let C_i , $i = 1, 2$ be irreducible curves of genus $g(C_i) \geq 2$. Then $(C_i, C_j) \geq 2$.*

Proof. If $C_1 - C_2$ or $C_2 - C_1$ is effective, then $(C_1, C_2) \geq (C_2^2) \geq 2$ or $(C_1, C_2) \geq (C_1^2) \geq 2$. If $C_1 - C_2$ and $C_2 - C_1$ are not effective, then $0 \geq \chi(\mathcal{O}_X(C_1 - C_2)) \geq (C_1 - C_2)^2/2$. Hence we see that $(C_1, C_2) \geq 2$. $\square$

Definition 3.4. For a Mukai vector $v \in H^{\text{ev}}(X, \mathbb{Z})$, $\mathcal{M}(v)$ is the stack of coherent sheaves E of $v(E) = v$.

Lemma 3.7. *Let E be a purely 1-dimensional sheaf such that $\text{Supp } E$ consists of genus $g \geq 2$ curves. Then $\dim \mathcal{M}(v(E)) = \langle v(E)^2 \rangle + 1$ at E , if H is general.*

Proof. We set $v := v(E)$. Let $\mathcal{M}$ be an irreducible component of $\mathcal{M}(v)$ containing E and let E' be a general point of $\mathcal{M}$. We consider the Harder-Narasimhan filtration of E' :

$$(3.19) \quad 0 \subset F_1 \subset F_2 \subset \cdots \subset F_s = E'.$$

We set $v_i = v(F_i/F_{i-1})$. By our assumption, we may assume that $\text{Supp } F_i/F_{i-1}$ consist of curves of genus greater than 1. Hence $\langle v_i^2 \rangle > 0$. Moreover by Lemma 3.6, $\langle v_i, v_j \rangle \geq 2$. Let $\mathcal{F}^0(v_1, v_2, \dots, v_s)$ be the stack of filtrations (3.19) such that $\text{Hom}(F_i/F_{i-1}, F_j/F_{j-1}) = 0$ for $i < j$. By [Y7, Lem. 5.2],

$$(3.20) \quad \dim \mathcal{F}^0(v_1, v_2, \dots, v_s) = \sum_{i=1}^s \dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} + \sum_{i < j} \langle v_i, v_j \rangle.$$

Since $\langle v_i^2 \rangle > 0$, we get $\dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} = \langle v_i^2 \rangle + 1$. Hence we see that

$$(3.21) \quad \begin{aligned} \dim \mathcal{F}^0(v_1, v_2, \dots, v_s) &= \langle v^2 \rangle + 1 - \left(\sum_{i=1}^s (\langle v_i \rangle + 1) + \sum_{i < j} \langle v_i, v_j \rangle \right) \\ &= \sum_{i < j} \langle v_i, v_j \rangle - (s-1) > 0. \end{aligned}$$

Since $\dim \mathcal{M}(v) \geq \langle v^2 \rangle + 1$, we get our claim. $\square$

Lemma 3.8. *Assume that X is a K3 surface. Let E be a purely 1-dimensional sheaf of $\text{Div}(E) = r\sigma$. Then $\mathcal{M}(v(E)) = -r^2$ at E .*

Proof. We set $v := v(E)$. Let $\mathcal{M}$ be an irreducible component of $\mathcal{M}(v)$ containing E and let E' be a general point of $\mathcal{M}$. We consider the Harder-Narasimhan filtration of E' :

$$(3.22) \quad 0 \subset F_1 \subset F_2 \subset \cdots \subset F_s = E'.$$

We set $v_i = v(F_i/F_{i-1})$. Then $v_i = r_i\sigma + a_i\omega_X$. It is easy to see that a_i is divisible by r_i and $\mathcal{M}_{\sigma+kf}(v_i)^{ss} = \{\mathcal{O}_\sigma(a_i/r_i - 1)^{\oplus r_i}\}$. Then $\dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} = -r_i^2$. As in Lemma 3.7, let $\mathcal{F}^0(v_1, v_2, \dots, v_s)$ be the stack of filtrations (3.22) such that $\text{Hom}(F_i/F_{i-1}, F_j/F_{j-1}) = 0$ for $i < j$. By [Y7, Lem. 5.2],

$$\begin{aligned} \dim \mathcal{F}^0(v_1, v_2, \dots, v_s) &= \sum_{i=1}^s \dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} + \sum_{i < j} \langle v_i, v_j \rangle \\ (3.23) \quad &= -\sum_{i=1}^s r_i^2 - \sum_{i < j} 2r_i r_j \\ &= -r^2. \end{aligned}$$

Therefore we get our claim. $\square$

Lemma 3.9. *Let E be a purely 1-dimensional sheaf on X such that $v(E) = rf + a\omega_X$, or $v(E) = r\sigma + a\omega_X$. Assume that $\langle v(E)^2 \rangle = 0$. Then $\dim \mathcal{M}(v(E)) = r$ at E .*

Proof. We set $v := v(E)$. Let $\mathcal{M}$ be an irreducible component of $\mathcal{M}(v)$ containing E and let E' be a general point of $\mathcal{M}$. We consider the Harder-Narasimhan filtration of E' :

$$(3.24) \quad 0 \subset F_1 \subset F_2 \subset \dots \subset F_s = E'.$$

We set $v_i = v(F_i/F_{i-1})$. Then we see that $\langle v_i^2 \rangle = 0$. As in the proof of Lemma 1.8 in [Y7], we see that $\dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} = r_i$. By using [Y7, Lem. 5.2] again, we see that

$$\begin{aligned} \dim \mathcal{F}^0(v_1, v_2, \dots, v_s) &= \sum_{i=1}^s \dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} + \sum_{i < j} \langle v_i, v_j \rangle \\ (3.25) \quad &= \sum_{i=1}^s r_i = r. \end{aligned}$$

Therefore we get our claim. $\square$

For N_1 , we get the following.

Proposition 3.10. $\dim N_1 < \dim \mathcal{M}_{\sigma+kf}(w)$.

Proof. Assume that $\text{Supp}(L)$ is not irreducible. Then there is a filtration

$$(3.26) \quad 0 \subset F_1 \subset F_2 \subset F_3 = L$$

such that (i) $\text{Div}(F_1) = r_1\sigma$ (ii) F_2/F_1 is a pure dimension 1 sheaf of $\text{Div}(F_2/F_1) = r_2f$ and (iii) F_3/F_2 is a pure dimension 1 sheaf of $\text{Div}(F_3/F_2) = C_3$, where C_3 consists of curves of genus greater than 1. We set $v_i := v(F_i/F_{i-1})$, $i = 1, 2, 3$. By Lemma 3.7, we may assume that $F_2 \neq 0$. We first note that $\text{Ext}^2(F_i/F_{i-1}, F_j/F_{j-1}) = \text{Hom}(F_j/F_{j-1}, F_i/F_{i-1})^\vee = 0$ for $i \neq j$.

(I) We first treat the case where X is a K3 surface. Assume that $F_2 \neq F_3$. By Lemmas 3.7, 3.8, 3.9, we see that

$$\begin{aligned} \dim \mathcal{F}^0(v_1, v_2, v_3) &= \sum_i \dim \mathcal{M}_{\sigma+kf}(v_i)^{ss} + \sum_{i < j} \langle v_i, v_j \rangle \\ (3.27) \quad &= \langle w^2 \rangle + 1 - (-r_1^2 - r_2 + (r_1\sigma, r_2f + C_3) + (r_2f, C_3)). \end{aligned}$$

By our assumption, $(c_1(w), \sigma) \geq 0$. Hence $(r_1\sigma, r_2f + C_3) \geq 2r_1^2$. By our assumption, $(f, C_3) > 0$. Therefore $-r_1^2 - r_2 + (r_1\sigma, r_2f + C_3) + (r_2f, C_3) \geq r_1^2 > 0$. We next assume that $F_2 = F_3$. Then we see that

$$(3.28) \quad \dim \mathcal{F}^0(v_1, v_2) = \langle w^2 \rangle + 1 - (-r_1^2 - r_2 + 1 + r_1r_2).$$

Since $(c_1(w), \sigma) \geq 0$, $r_2 \geq 2r_1$. Then $-r_1^2 - r_2 + 1 + r_1r_2 \geq r_1(r_1 - 2) + 1 \geq 1$, because $c_1(w)$ is not primitive. Therefore we get our claim.

(II) We next treat the case where X is an abelian surface. Assume that $F_2 \neq F_3$. Then

$$(3.29) \quad \dim \mathcal{F}^0(v_1, v_2, v_3) = \langle w^2 \rangle + 1 - (-r_1 - r_2 + (r_1\sigma, r_2f + C_3) + (r_2f, C_3)).$$

Since C_3 consists of curves of genus greater than 1, $(\sigma, C_3) > 1$ and $(f, C_3) > 1$. Then $(-r_1 - r_2 + (r_1\sigma, r_2f + C_3) + (r_2f, C_3)) > 0$. If $F_2 = F_3$, then

$$(3.30) \quad \dim \mathcal{F}^0(v_1, v_2) = \langle w^2 \rangle + 1 - (-r_1 - r_2 + 1 + r_1r_2).$$

Since $c_1(w)$ is not primitive, $(r_1 - 1)(r_2 - 1) > 0$. Therefore we get our claim. $\square$

Estimate of $\dim N_2$. For an integer $\lambda \geq 2$, let $\text{Hilb}_X^\xi(\lambda)$ be the locally closed subset of Hilb_X^ξ consisting of λC , where C is an integral curve. By Proposition 3.10, it is sufficient to estimate the dimension of $\text{Hilb}_X^\xi(\lambda)$. Let C be an integral curve. For $D = \lambda C$ and $w = D + r\omega_X$, we set

$$(3.31) \quad M_{\sigma+kf}(w, D) := \{L \in M_{\sigma+kf}(w) \mid \text{Div}(L) = D\}.$$

We fix a point $x \in C$. We also set

$$(3.32) \quad P_{\sigma+kf}(w, w + \omega_X, D) := \left\{ L \subset L' \left| \begin{array}{l} L' \in M_{\sigma+kf}(w + \omega_X), \text{Div}(L') = D \\ L \in M_{\sigma+kf}(w), L'/L \cong \mathbb{C}_x \end{array} \right. \right\}.$$

Let π_w and $\pi_{w+\omega_X}$ be natural projections sending $L \subset L'$ to L and L' respectively:

$$(3.33) \quad \begin{array}{ccc} & P_{\sigma+kf}(w, w + \omega_X, D) & \\ \pi_w \swarrow & & \searrow \pi_{w+\omega_X} \\ M_{\sigma+kf}(w, D) & & M_{\sigma+kf}(w + \omega_X, D) \end{array}$$

For $L \in M_{\sigma+kf}(w, D)$, $-\dim \text{Ext}^1(\mathbb{C}_x, L) + \dim \text{Ext}^2(\mathbb{C}_x, L) = \chi(\mathbb{C}_x, L) = 0$. Combining Serre duality, we see that $\dim \text{Ext}^1(\mathbb{C}_x, L) = \dim \text{Ext}^2(\mathbb{C}_x, L) = \dim \text{Hom}(L, \mathbb{C}_x)$. By the following lemma, $\dim \pi_w^{-1}(L) \leq \lambda - 1$ and $\dim \pi_{w+\omega_X}^{-1}(L') \leq \lambda - 1$.

Lemma 3.11. *Let x be a smooth point of C . Let L be a purely 1-dimensional sheaf such that $\text{Div}(L) = \lambda C$. Then $\dim L \otimes \mathbb{C}_x \leq \lambda$.*

Proof. Let C' be a germ of a curve intersecting C at x transversely. Let $\mathcal{O}_{X,x}$ be the stalk of $\mathcal{O}_X$ at x . We take a free resolution of $L \otimes \mathcal{O}_{X,x}$:

$$(3.34) \quad 0 \rightarrow \mathcal{O}_{X,x}^{\oplus n} \xrightarrow{A} \mathcal{O}_{X,x}^{\oplus n} \rightarrow L \otimes \mathcal{O}_{X,x} \rightarrow 0.$$

Then the local equation of $\text{Div}(L)$ at x is given by $\det(A)$. By restricting the sequence to C' , we get a free resolution of $L \otimes \mathcal{O}_{C',x}$. Then $\dim(L \otimes \mathcal{O}_{C',x})$ is given by the local intersection number $(\text{Div}(D), C')_x = \lambda$. Therefore we get our claim. $\square$

Lemma 3.12. $\dim M_{\sigma+kf}(w + \omega_X, D) = (D^2)/2 + 1$.

Proof. By [Y8, Thm. 3.15], $M_{\sigma+kf}(w + \omega_X)$ is isomorphic to $M_{\sigma+kf}(r + f - r\omega_X)$. Since $r + f - r\omega_X$ is primitive, [Y8, Thm. 0.1 and 8.1] implies that it is irreducible. For a smooth curve $C \in \text{Hilb}_X^\xi$, the fiber of $M_{\sigma+kf}(w + \omega_X) \rightarrow \text{Hilb}_X^\xi$ is $\text{Pic}^{r+1}(C)$. It is easy to see that $\text{Pic}^{r+1}(C)$ is a Lagrangian subscheme of $M_{\sigma+kf}(w + \omega_X)$. By Matsushita [Mt], every fiber is of dimension $(\xi^2)/2 + 1$. $\square$

Lemma 3.13. *Let L be a stable sheaf of $v(L) = w$ and $\text{Div}(L) = D$, and let L' be a coherent sheaf which fits in a non-trivial extension*

$$(3.35) \quad 0 \rightarrow L \rightarrow L' \rightarrow \mathbb{C}_x \rightarrow 0$$

where $x \in D$. Then L' is stable.

Proof. Assume that L' is not of pure dimension 1 and let T be the 0-dimensional subsheaf of L' . Then $T \rightarrow L' \rightarrow \mathbb{C}_x$ must be injective. Hence it is isomorphic, which implies that the exact sequence split. Therefore L' is of pure dimension 1. If L' is not stable, then there is a subsheaf L_1 of L' such that $\frac{\chi(L_1)}{\lambda_1} > \frac{\chi(L')}{\lambda}$, where $\text{Div}(L_1) = \lambda_1 C$. Hence $0 < \chi(L_1)\lambda - \chi(L')\lambda_1 = (\chi(L_1) - 1)\lambda - \chi(L)\lambda_1 + \lambda - \lambda_1$. Since $\chi(L) = r$ is divisible by λ and $\lambda - \lambda_1 < \lambda$, we get $(\chi(L_1) - 1)\lambda - \chi(L)\lambda_1 \geq 0$. Thus $\frac{\chi(L_1 \cap L)}{\lambda_1} \geq \frac{\chi(L)}{\lambda}$, which implies that L is not stable. Therefore L' must be stable. $\square$

Corollary 3.14. $\dim M_{\sigma+kf}(w, D) \leq \dim P_{\sigma+kf}(w, w + \omega_X, D) \leq \dim M_{\sigma+kf}(w + \omega_X, D) + (\lambda - 1)$.

Proof. By Lemma 3.13, π_w is surjective. Hence we get our claim from Lemma 3.12. $\square$

Since $\lambda \geq 2$, we get that $2\lambda^2 - (\lambda^2 + \lambda + 1) > 0$. Then

$$(3.36) \quad \begin{aligned} \dim j^{-1}(\text{Hilb}_X^\xi(\lambda)) &\leq (C^2)/2 + 1 + \lambda^2(C^2)/2 + 1 + (\lambda - 1) \\ &< (\lambda^2 + \lambda + 1)(C^2)/2 + 2 \\ &< \lambda^2(C^2) + 2 = \dim M_{\sigma+kf}(w). \end{aligned}$$

Combining Proposition 3.10, we get the following.

Proposition 3.15. *We set*

$$(3.37) \quad \begin{aligned} M_{\sigma+kf}(w)_0 &:= \{L \in M_{\sigma+kf}(w) \mid \text{Div}(L) \text{ is an integral curve} \} \\ &= M_{\sigma+kf}(w) \setminus (N_1 \cup N_2). \end{aligned}$$

Then $M_{\sigma+kf}(w)_0$ is an open dense subscheme of $M_{\sigma+kf}(w)$.

Proposition 3.16. *$M_{\sigma+kf}(w)$ is irreducible.*

Proof. By Proposition 3.15, it is sufficient to show that $M_{\sigma+kf}(w)_0$ is irreducible. Let C be an integral curve. Then $j^{-1}(C)$ is the compactified Jacobian of C . By [A-I-K], the compactified Jacobian of C is irreducible. Therefore $M_{\sigma+kf}(w)_0$ is irreducible. $\square$

Combining all together, we get the following theorem.

Theorem 3.17. *Let X be an abelian surface or a K3 surface and let v be a Mukai vector of $\text{rk } v > 0$ and $\langle v^2 \rangle > 0$. Let H be a general ample divisor with respect to v . Then $\mathcal{M}_H(v)^{ss}$ is a normal and irreducible stack. In particular, $\overline{\mathcal{M}}_H(v)$ is a normal variety.*

Remark 3.4. In [O2], [O3], O'Grady studied the case where $m(v) = 2$. In particular, he constructed symplectic desingularization of $\overline{\mathcal{M}}_H(v)$, if $\langle v^2 \rangle = 8$.

4. FOURIER-MUKAI TRANSFORM ON ENRIQUES SURFACES

In this section, we consider Fourier-Mukai transform on Enriques surface X . By using Fourier-Mukai transform, we shall compute Hodge polynomial of some moduli spaces of sheaves.

In our case, Mukai vector $v(x)$ of $x \in K(X)$ is defined as an element of $H^*(X, \mathbb{Q})$:

$$(4.1) \quad \begin{aligned} v(x) &:= \text{ch}(x) \sqrt{\text{td}_X} \\ &= \text{rk}(x) + c_1(x) + \left(\frac{\text{rk}(x)}{2} \omega_X + \text{ch}_2(x)\right) \in H^*(X, \mathbb{Q}). \end{aligned}$$

We also introduce Mukai's pairing on $H^*(X, \mathbb{Q})$ by $\langle x, y \rangle := -\int_X x^\vee \wedge y$. Then we have an isomorphism of lattices:

$$(4.2) \quad (v(K(X)), \langle \cdot, \cdot \rangle) \cong \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus E_8(-1).$$

Definition 4.1. We call an element of $v(K(X))$ by Mukai vector. A Mukai vector v is primitive, if v is primitive as an element of $v(K(X))$.

The following was essentially proved in [Y1, sect. 3.2].

Proposition 4.1. *Let X be a surface such that K_X is numerically trivial. Let (H, α) be a pair of ample divisor H and a $\mathbb{Q}$ -divisor α . Then $e(\mathcal{M}_H^\alpha(v)^{ss})$ does not depend on the choice of H and α , if $(H, \mathcal{O}(\alpha))$ is general with respect to v (cf. Defn. 1.5).*

Proposition 4.2. *Let v be a Mukai vector such that $\text{rk}(v)$ is odd. Then $\mathcal{M}_H(v)^s$ is smooth of $\dim \mathcal{M}_H(v)^s = \langle v^2 \rangle + 1$.*

Proof. For $E \in \mathcal{M}_H(v)^s$, we get $\det(E(K_X)) \not\cong \det(E)$. If there is a non-zero homomorphism $E \rightarrow E(K_X)$, then the stability implies that it is an isomorphism. Hence $\text{Ext}^2(E, E) = \text{Hom}(E, E(K_X))^\vee = 0$. Since $-\chi(E, E) = -\langle v(E), v(E) \rangle$, $\mathcal{M}_H(v)^s$ is smooth of $\dim \mathcal{M}_H(v)^s = \langle v^2 \rangle + 1$. $\square$

For a Mukai vector v , let $L_1, L_2 = L_1(K_X) \in \text{Pic}(X)$ be line bundles on X such that $c_1(L_1) (= c_1(L_2)) = c_1(v)$. Then we have a decomposition

$$(4.3) \quad \mathcal{M}_H(v)^{ss} = \mathcal{M}_H(v, L_1)^{ss} \coprod \mathcal{M}_H(v, L_2)^{ss}$$

where $\mathcal{M}_H(v, L_i)^{ss}$, $i = 1, 2$, is the substack of $\mathcal{M}_H(v)^{ss}$ consisting of E such that $\det(E) = L_i$. We also have a decomposition $M_H(v) = M_H(v, L_1) \coprod M_H(v, L_2)$, where $M_H(v, L_i)$ is the subscheme of $M_H(v)$ consisting of E such that $\det(E) = L_i$.

We consider Fourier-Mukai transform associated to (-1) -reflection. Let $v_0 := r + c_1 - (s/2)\omega_X$ be a Mukai vector such that $\text{rk}(v_0) > 0$ and $\langle v_0^2 \rangle = (c_1^2) + rs = -1$. Since (c_1^2) is even, r and s are odd. Let H be a general ample divisor with respect to v_0 . Assume that there is a stable vector bundle E_0 with respect to H such that $v(E_0) = v_0$ (cf. Corollary 4.7). Then we see that

$$(4.4) \quad \begin{cases} \text{Hom}(E_0, E_0) = \mathbb{C} \\ \text{Ext}^1(E_0, E_0) = 0 \\ \text{Ext}^2(E_0, E_0) = 0. \end{cases}$$

Let

$$(4.5) \quad \begin{aligned} ev_1 : E_0^\vee \boxtimes E_0 &\rightarrow \mathcal{O}_\Delta, \\ ev_2 : E_0(K_X)^\vee \boxtimes E_0(K_X) &\rightarrow \mathcal{O}_\Delta \end{aligned}$$

be evaluation maps. We define a sheaf $\mathcal{E}$ on $X \times X$ by an exact sequence

$$(4.6) \quad 0 \rightarrow \mathcal{E} \rightarrow E_0^\vee \boxtimes E_0 \oplus E_0(K_X)^\vee \boxtimes E_0(K_X) \xrightarrow{(ev_1, ev_2)} \mathcal{O}_\Delta \rightarrow 0.$$

Then $\mathcal{E}_{|\{x\} \times X}$ (resp. $\mathcal{E}_{|X \times \{x\}}$) is a stable sheaf of $v(\mathcal{E}_{|\{x\} \times X}) = 2 \operatorname{rk}(E_0)v(E_0) - \omega_X$ (resp. $v(\mathcal{E}_{|X \times \{x\}}) = 2 \operatorname{rk}(E_0)v(E_0)^\vee - \omega_X$). Thus $\mathcal{E}$ is a flat family of stable sheaves of $v(\mathcal{E}_{|\{x\} \times X}) = 2 \operatorname{rk}(E_0)v(E_0) - \omega_X$. By the construction of $\mathcal{E}$, $\mathcal{E}_{|\{x\} \times X}(K_X) \cong \mathcal{E}_{|\{x\} \times X}$, which implies that

$$(4.7) \quad \operatorname{Ext}^2(\mathcal{E}_{|\{x\} \times X}, \mathcal{E}_{|\{x\} \times X}) = \operatorname{Hom}(\mathcal{E}_{|\{x\} \times X}, \mathcal{E}_{|\{x\} \times X}(K_X))^\vee \cong \mathbb{C}.$$

Since $\langle v(E_0)^2 \rangle = -1$, we see that $\langle v(\mathcal{E}_{|\{x\} \times X})^2 \rangle = 0$. Hence the Zariski tangent space is 2-dimensional:

$$(4.8) \quad \operatorname{Ext}^1(\mathcal{E}_{|\{x\} \times X}, \mathcal{E}_{|\{x\} \times X}) \cong \mathbb{C}^{\oplus 2}.$$

Therefore X is a connected component of $M_H(v_1)$, where $v_1 = 2 \operatorname{rk}(E_0)v(E_0) - \omega_X$.

Then $\mathcal{H}_\mathcal{E} : \mathbf{D}(X) \rightarrow \mathbf{D}(X)_{op}$ is an equivalence of categories. As a corollary of this fact, we get that $M_H(v_1) = X$. By our construction of $\mathcal{E}$, we see that

$$(4.9) \quad v(\mathcal{H}_\mathcal{E}(x)) = -(x^\vee + 2v(E_0)^\vee \langle x, v(E_0) \rangle).$$

If $E_0 = \mathcal{O}_X$ and $v(E) = r + c_1 + (s/2)\omega_X$, then $v(\mathcal{H}_\mathcal{E}(E)) = s + c_1 + (r/2)\omega_X$.

From now on, we assume that X is unnodal, i.e. there is no (-2) -curve. Let σ and f be elliptic curves on X such that $(\sigma, f) = 1$. Then

$$(4.10) \quad H^2(X, \mathbb{Z})_f = (\mathbb{Z}\sigma \oplus \mathbb{Z}f) \perp E_8(-1)$$

where $H^2(X, \mathbb{Z})_f$ is the torsion free quotient of $H^2(X, \mathbb{Z})$.

Proposition 4.3. *We set $G_1 := \mathcal{E}_{|\{x\} \times X}$ and $G_2 := \mathcal{E}_{|X \times \{x\}}$. Assume that $\deg_{G_1}(v) = 0$ and $l(v) := -\langle v, \omega_X \rangle / \operatorname{rk} v_1 > 0$, $a(v) := \langle v, v_1 \rangle / \operatorname{rk} v_1 > 0$. Let ε be an element of $K(X) \otimes \mathbb{Q}$ such that $v(\varepsilon) \in v_1^\perp \cap \omega_X^\perp$, $|\langle v(\varepsilon)^2 \rangle| \ll 1$ and $(H, c_1(\varepsilon)) = 0$. Then $\mathcal{H}_\mathcal{E}$ induces an isomorphism*

$$(4.11) \quad \mathcal{M}_H^{G_1+\varepsilon}(v)^{ss} \rightarrow \mathcal{M}_H^{G_2+\hat{\varepsilon}}(-\mathcal{H}_\mathcal{E}(v))^{ss}.$$

Proof. Since H is general with respect to $v(E_0)$, we see that $\mathcal{E}_{|\{x\} \times X}$ is G_1 -twisted stable. Then we see that Lemma 2.4 holds. We next show that Lemma 2.5 holds. We may assume that E is μ -stable. If $E^{\vee\vee} \neq E_0, E_0(K_X)$, then $\operatorname{Hom}(E, E_0) = \operatorname{Hom}(E, E_0(K_X)) = 0$. If $E^{\vee\vee} = E_0, E_0(K_X)$, then $\operatorname{Hom}(E, \mathcal{E}_{|\{x\} \times X}) = 0$ for $x \in X \setminus \operatorname{Supp}(E^\vee/E)$. Thus Lemma 2.5 holds. Then the same proof of Theorem 2.3 works and we get our claim. $\square$

Corollary 4.4. $M_H(r - (1/2)\omega_X, \mathcal{O}_X) \cong \operatorname{Hilb}_X^{(r+1)/2}$ for a general H with respect to $r - (1/2)\omega_X$.

Proposition 4.5. *Assume that $r, s > 0$. Then $e(M_H^\alpha(r + c_1 - (s/2)\omega_X)) = e(M_H^\alpha(s - c_1 - (r/2)\omega_X))$ for a general (H, α) , if $\langle c_1^2 \rangle < 0$, i.e. $\langle v^2 \rangle < rs$, where $v = r + c_1 - (s/2)\omega_X$. In particular, if $r > \langle v^2 \rangle$, then we get our claim.*

Proof. If $\langle c_1^2 \rangle < 0$, then Hodge index theorem implies that there is a divisor H such that $(H, c_1) = 0$ and $\langle H^2 \rangle > 0$. By Riemann-Roch theorem, we may assume that H is effective. Since X is unnodal, H is ample. If $E_0 = \mathcal{O}_X$, then $v(\mathcal{E}_{|\{x\} \times X}) = 2$. Hence v satisfies assumptions of Proposition 4.3. Then we get an isomorphism

$$(4.12) \quad M_H^{\mathcal{O}_X+\varepsilon}(r + c_1 - (s/2)\omega_X) \rightarrow M_H^{\mathcal{O}_X+\varepsilon}(s - c_1 - (r/2)\omega_X),$$

where $(H, \mathcal{O}_X + \varepsilon)$ is general with respect to v . By Proposition 4.1, we get our claim. $\square$

Theorem 4.6. *Let $v = r + c_1 - (s/2)\omega_X \in H^*(X, \mathbb{Q})$ be a primitive Mukai vector such that r is odd. Then*

$$(4.13) \quad e(M_H(v, L)) = e(\operatorname{Hilb}_X^{(\langle v^2 \rangle + 1)/2})$$

for a general H , where $L \in \operatorname{Pic}(X)$ satisfies $c_1(L) = c_1$. In particular,

- (i) $M_H(v) \neq \emptyset$ for a general H if and only if $\langle v^2 \rangle \geq -1$.
- (ii) $M_H(v, L)$ is irreducible for a general H .

Proof. We first assume that $c_1 \in E_8(-1)$. We set $l = \gcd(r, c_1)$. Replacing v by $v \exp(\xi_1)$, $\xi_1 \in E_8(-1)$, we may assume that c_1/l is primitive and $s > \langle v^2 \rangle$. Since v is primitive, $\gcd(l, s) = 1$. By Proposition 4.5, we get

$$(4.14) \quad e(M_H(r + c_1 - (s/2)\omega_X)) = e(M_H(s - c_1 - (r/2)\omega_X)).$$

Replacing $v = r + c_1 - (s/2)\omega_X$ by $v' = s - c_1 - (r/2)\omega_X$, we may assume that $r > \langle v^2 \rangle$. By the same argument as above, we may assume that $l = 1$ and c_1 is primitive. We set $D = \sigma - \frac{(\eta^2)}{2}f + \eta$, where $\eta \in E_8(-1)$ satisfies that $2(\eta, c_1) = s - 1$. Then $(D^2) = 0$ and $-s + 2(c_1, D) = -1$. Since $v \exp(D) = r + (c_1 + rD) - 1/2\omega_X$, $e(M_H(v)) = e(M_H(r + (c_1 + rD) - 1/2\omega_X))$. Since $r > \langle v^2 \rangle$, Proposition 4.5 implies that our claim holds for this case.

We shall next treat the general case. We use induction on r . We set $c_1 := d_1\sigma + d_2f + \xi$. Replacing v by $v \exp(k\sigma)$, we may assume that $0 \leq |d_1| < r/2$. We first assume that $d_1 \neq 0$. We note that $(c_1, f) = d_1$. Replacing v by $v \exp(\eta)$, $\eta \in E_8(-1)$, we may assume that $s > \langle v^2 \rangle$. Then by Proposition 4.5, $e(M_H(v)) = e(M_H(s - c_1 - (r/2)\omega_X))$ for a general H . We take an integer k such that $0 < r + 2d_1k < 2|d_1| < r$. Then $v \exp(kf) = s + (-c_1 + skf) - r'/2\omega_X$, where $r' = r + 2d_1k$. Since $s > \langle v^2 \rangle$, Proposition 4.5, implies that $e(M_H(s + (-c_1 + skf) - r'/2\omega_X)) = e(M_H(r' + (c_1 - skf) - (s/2)\omega_X))$ for a general H . By induction hypothesis, we get our claim.

If $d_1 = 0$, then we may assume that $0 \leq |d_2| < r/2$. If $|d_2| > 0$, then we can apply the same argument and get our claim. If $d_1 = d_2 = 0$, then $c_1 \in E_8(-1)$, so we get our claim. $\square$

Corollary 4.7. *If $\langle v^2 \rangle = -1$, then there is a stable vector bundle E_0 of $v(E_0) = v$ with respect to H .*

Remark 4.1. By the proof, we also get the following: Let v be a primitive Mukai vector such that $\text{rk } v$ is odd. Then $e(\mathcal{M}_H(mv)^{ss}) = e(\mathcal{M}_H(m(1 - (n/2)\omega_X))^{ss})$, where $n = \langle v^2 \rangle$.

4.1. Relation to Montonen-Olive duality. We fix an Enriques surface X and fix the following data:

- $(H^2(X, \mathbb{Z})_f, Q) : a$ lattice with $Q(x, y) = -\int_X x \wedge y$, $x, y \in H^2(X, \mathbb{Z})_f$.
- A orthogonal decomposition of $H^2(X, \mathbb{R})$ as a sum of definite signature:

$$(4.15) \quad P : H^2(X, \mathbb{Z}) \otimes \mathbb{R} \cong \mathbb{R}^{9,0} \oplus \mathbb{R}^{0,1}.$$

- $P_L(x) = x_L$, $P_R(x) = x_R$: the projections onto the two factors.

For an odd integer $r > 0$, we define $U(r)$ -partition function by

$$(4.16) \quad Z^r(\tau, x) := \sum_{\substack{v \in H^*(X, \mathbb{Q}) \\ \text{rk } v = r}} " \chi(M_H(v)) " q^{\frac{\langle v^2 \rangle}{2r}} q^{\frac{1}{2r}Q(c_1(v)_L^2)} \bar{q}^{\frac{-1}{2r}Q(c_1(v)_R^2)} e^{Q(c_1(v), x)},$$

where $(\tau, x) \in \mathbb{H} \times H^2(X, \mathbb{Z}) \otimes \mathbb{C}$, $\mathbb{H} := \{\tau \in \mathbb{C} | \Im \tau > 0\}$, $q := \exp(2\pi\sqrt{-1}\tau)$, $e := \exp(2\pi\sqrt{-1})$ and " $\chi(M_H(v))$ " is a kind of Euler characteristic of a nice compactification of $M_H(v)$ (Mathematically there is no definition). Then Montonen-Olive duality for $U(r)$ gauge group (cf. [V-W]) says that

(#) $Z^r(\tau, x)$ transforms like a Jacobi form of holomorphic/anti-holomorphic weight

$$(4.17) \quad (-\chi(X)/2 + b_-(X)/2, b_+(X)/2) = (-3/2, 1/2).$$

$SU(1)$ -partition function $Z_0^1(\tau)$ is given by $Z_0^1(\tau) = 1/\eta(\tau)^{12}$ ([Gö], [V-W]). Hence

$$(4.18) \quad \begin{aligned} Z^1(\tau, x) &= 2Z_0^1(\tau) \left(\sum_{c \in H^2(X, \mathbb{Z})_f} q^{\frac{1}{2}Q(c_L^2)} \bar{q}^{\frac{-1}{2}Q(c_R^2)} e^{Q(c, x)} \right) \\ &= \frac{2}{\eta(\tau)^{12}} \Theta(\tau, P, x), \end{aligned}$$

where $\Theta(\tau, P, x) = \sum_{c \in H^2(X, \mathbb{Z})_f} q^{\frac{1}{2}Q(c_L^2)} \bar{q}^{\frac{-1}{2}Q(c_R^2)} e^{Q(c, x)}$ and the factor 2 comes from the torsion submodule of $H^2(X, \mathbb{Z})$. We set

$$(4.19) \quad \Gamma := \left\langle \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\rangle \subset SL_2(\mathbb{Z}).$$

For an odd integer $r > 0$, we set

$$(4.20) \quad N(r) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \pmod{2} \right\}.$$

Then we have a coset decomposition:

$$(4.21) \quad N(r) = \prod_{\substack{a,b,d>0 \\ ad=r, 0 \leq b < d}} \Gamma \begin{pmatrix} a & 2b \\ 0 & d \end{pmatrix}.$$

Since $\Theta(\tau, P, x)$ transforms like a Jacobi form of holomorphic/anti-holomorphic weight $(9/2, 1/2)$, $Z^1(\tau, x)$ transforms like a Jacobi form of holomorphic/anti-holomorphic weight $(-3/2, 1/2)$:

$$(4.22) \quad Z^1 \left(-\frac{1}{\tau}, \frac{x_L}{\tau} + \frac{x_R}{\bar{\tau}} \right) = (-\sqrt{-1}\tau)^{-3/2} (\sqrt{-1}\bar{\tau})^{1/2} e^{\frac{Q(x_L^2)}{2\tau}} e^{\frac{Q(x_R^2)}{2\bar{\tau}}} Z^1(\tau, x_L + x_R).$$

We conjecture that $Z^r(\tau, x)$ is given by Hecke transformation of order r of $Z^1(\tau, x)$ ([MNVW]):

$$(4.23) \quad Z^r(\tau, x) = \frac{1}{r^2} \sum_{\substack{a,b,d \geq 0 \\ ad=r, b < d}} dZ^1 \left(\frac{a\tau + 2b}{d}, ax \right).$$

Hence $Z^r(\tau, x)$ transforms like a Jacobi form of holomorphic/anti-holomorphic weight $(-3/2, 1/2)$ and index r . Thus (#) holds.

Evidence:

We shall give an evidence of this conjecture by using Theorem 4.6. For simplicity, we set $X^{[n]} = \text{Hilb}_X^n$. Then we see that

$$(4.24) \quad \begin{aligned} & \frac{1}{2} \sum_{0 \leq b < d} dZ^1 \left(\frac{a\tau + 2b}{d}, ax \right) \\ &= \sum_{0 \leq b < d} \sum_{\xi \in H^2(X, \mathbb{Z})_f} \sum_n d\chi(X^{[n]}) q^{\frac{a}{d}(n-1/2)} q^{\frac{a}{2d}Q(\xi_L^2)} \bar{q}^{\frac{-a}{2d}Q(\xi_R^2)} e^{aQ(\xi, x)} e^{\frac{2b}{d}((n-1/2)+Q(\xi^2)/2)} \\ &= \sum_{\xi \in H^2(X, \mathbb{Z})_f} \sum_{d|2n-1+Q(\xi^2)} d^2 \chi(X^{[n]}) q^{\frac{a}{d}(n-1/2)} q^{\frac{a}{2d}Q(\xi_L^2)} \bar{q}^{\frac{-a}{2d}Q(\xi_R^2)} e^{aQ(\xi, x)} \\ &= \sum_{\xi \in H^2(X, \mathbb{Z})_f} \sum_k d^2 \chi(X^{[(kd+1)/2-Q(\xi^2)/2]}) q^{\frac{a}{d}(kd/2-Q(\xi^2)/2)} q^{\frac{a}{2d}Q(\xi_L^2)} \bar{q}^{\frac{-a}{2d}Q(\xi_R^2)} e^{aQ(\xi, x)} \\ &= \sum_{\xi \in H^2(X, \mathbb{Z})_f} \sum_{w=(d, \xi, -k/2)} d^2 \chi(X^{[(\langle w^2 \rangle + 1)/2]}) q^{\frac{a}{2d}\langle w^2 \rangle} q^{\frac{a}{2d}Q(c_1(w)_L^2)} \bar{q}^{\frac{-a}{2d}Q(c_1(w)_R^2)} e^{aQ(c_1(w), x)} \\ &= \sum_{\text{rk } w=d} d^2 \chi(X^{[(\langle w^2 \rangle + 1)/2]}) q^{\frac{1}{2r}\langle (aw)^2 \rangle} q^{\frac{1}{2r}Q(c_1(aw)_L^2)} \bar{q}^{\frac{-1}{2r}Q(c_1(aw)_R^2)} e^{Q(c_1(aw), x)}. \end{aligned}$$

Therefore we get

$$(4.25) \quad \chi(M_H(v)) = \sum_{v=aw} \frac{2}{a^2} \chi(X^{[(\langle w^2 \rangle + 1)/2]}).$$

If v is primitive, by Theorem 4.6, we get

$$(4.26) \quad \chi(M_H(v)) = \chi(X^{[(\langle v^2 \rangle + 1)/2]}) = \chi(M_H(v)).$$

This implies that $\chi(M_H(v))$ is related to modular forms and in particular Hecke transforms.

5. TWISTED STABILITY FOR DIMENSION 1 SHEAVES

We shall generalize twisted stability to purely 1-dimensional sheaves. Let X be a smooth projective surface as in section 1.

Definition 5.1. Let E be a purely 1-dimensional sheaf on X .

(i) For a $\mathbb{Q}$ -divisor α on X (or its numerical equivalence class), we define α -twisted slope of E by

$$(5.1) \quad \mu_\alpha(E) := \frac{\chi(E(-\alpha))}{(c_1(E), H)} = \frac{\chi(E) - (c_1(E), \alpha)}{(c_1(E), H)}.$$

(ii) E is α -twisted semi-stable (resp. stable) if

$$(5.2) \quad \mu_\alpha(F) \leq_{(<)} \mu_\alpha(E)$$

for all $0 \subsetneq F \subsetneq E$.

(iii) For a vector bundle G on X (or its class in $K(X)$), we also define $\mu_G(E)$ by

$$(5.3) \quad \mu_G(E) := \frac{\chi_G(E)}{\deg_G(E)} = \frac{\chi(E) - (c_1(E), c_1(G))/\text{rk } G}{(c_1(E), H)} = \mu_{c_1(G)/\text{rk } G}(E).$$

We also define G -twisted stability as $c_1(G)/\text{rk } G$ -twisted stability.

Definition 5.2. Let $\mathcal{M}_H^\alpha(\gamma)^{ss}$ be the moduli stack of α -twisted semi-stable sheaves E of $\gamma(E) = \gamma$ and $\mathcal{M}_H^\alpha(\gamma)^s$ the open substack consisting of α -twisted stable sheaves.

We next generalize wall and chamber structure. Let $\mathcal{D}_\xi$ be a set of effective divisors D such that there is an effective divisor D' and $D + D' \equiv \xi$. It is known that $\mathcal{D}_\xi$ is a bounded set. For $E \in K(X)$ of $\gamma(E) = (0, \xi, \chi)$, we consider the set $\mathcal{W}$ of $\alpha \in \text{NS}(X) \otimes \mathbb{Q}$ which satisfies

$$(5.4) \quad \frac{\chi - (\xi, \alpha)}{(\xi, H)} = \frac{n - (D, \alpha)}{(D, H)},$$

for some $n \in \mathbb{Z}$, $D \in \mathcal{D}_\xi$ and $(D, n) \notin \mathbb{Q}(\xi, \chi)$. This set is a countable union of hyperplanes of $\text{NS}(X) \otimes \mathbb{Q}$. We claim that the numbers of hyperplanes is locally finite. Proof of the claim: Assume that α belongs to a bounded subset of $\text{Pic}(X) \otimes \mathbb{Q}$. Since the choice of D is finite, the set of n is bounded, which implies that the choice of n is finite.

Definition 5.3. We call a defining hyperplane of $\mathcal{W}$ by wall and a connected component of $\text{NS}(X) \otimes \mathbb{R} \setminus \mathcal{W}$ by chamber.

For α, α' which belong to a chamber $\mathcal{C}$, $\mathcal{M}_H^\alpha(\gamma)^{ss} = \mathcal{M}_H^{\alpha'}(\gamma)^{ss}$. Hence we may denote this stack by $\mathcal{M}_H^\mathcal{C}(\gamma)^{ss}$.

Definition 5.4. Let W be a wall and $\mathcal{C}$ a chamber such that $\overline{\mathcal{C}}$ intersects W . Let α be a $\mathbb{Q}$ -divisor belonging to $\overline{\mathcal{C}} \cap W$ and α_1 a $\mathbb{Q}$ -divisor which belongs to $\mathcal{C}$. $V^{\alpha, \mathcal{C}}(\gamma)$ be the set of α -twisted semi-stable sheaves with respect to H such that E is not α_1 -twisted semi-stable with respect to H and $\gamma(E) = \gamma$.

In the same way as in [Y1, Prop. 2.4], we get the following.

Proposition 5.1. Let C be a 2-dimensional vector space such that $C \cap \mathcal{C} \neq \emptyset$ and $\alpha \in C$.

(1) There is an element $\alpha_1 \in C$ such that $V^{\alpha, \mathcal{C}}(\gamma)$ is the set of torsion free sheaves E whose Harder-Narasimhan filtration

$$(5.5) \quad 0 \subset F_1 \subset F_2 \subset \cdots \subset F_s = E$$

with respect to α_1 satisfies $\mu_\alpha(F_i) = \mu_\alpha(F_{i+1})$ and is also the Harder-Narasimhan filtration of E with respect to $\alpha_t := \alpha_1 + t(\alpha - \alpha_1)$, $0 \leq t < 1$.

$$(2) \mathcal{M}_H^\alpha(\gamma)^{ss} = \mathcal{M}_H^\mathcal{C}(\gamma)^{ss} \amalg V^{\alpha, \mathcal{C}}(\gamma).$$

Lemma 5.2. Assume that $-K_X$ is base point free. Let E and F be α -twisted semi-stable sheaves of $\mu_\alpha(E) > \mu_\alpha(F)$, or of $\mu_\alpha(E) = \mu_\alpha(F)$ and $(c_1(F), -K_X) > 0$. Then $\text{Hom}(E, F(K_X)) = 0$.

Proof. Considering Jordan-Hölder filtration, we may assume that E and F are α -twisted stable. Since $-K_X$ is base point free, there is an inclusion $F(K_X) \hookrightarrow F$. Hence we get an inclusion $\text{Hom}(E, F(K_X)) \hookrightarrow \text{Hom}(E, F)$. If $\mu_\alpha(E) > \mu_\alpha(F)$, then $\text{Hom}(E, F) = 0$, and hence $\text{Hom}(E, F(K_X)) = 0$. Assume that $\mu_\alpha(E) = \mu_\alpha(F)$. If $\text{Hom}(E, F) \neq 0$, then $E \cong F$ and $\text{Hom}(E, F) \cong \mathbb{C}$. If there is a map $E \rightarrow F(K_X)$, then $E \rightarrow F(K_X) \subset F$ is not an isomorphism, because of $(c_1(F), -K_X) > 0$. Hence $\text{Hom}(E, F(K_X)) = 0$. $\square$

Corollary 5.3. If $-K_X$ is base point free and $(c_1(x), -K_X) > 0$, then $\mathcal{M}_H^\alpha(\gamma(x))^{ss}$ is smooth.

Proposition 5.4. Let C be a 2-dimensional vector space such that $C \cap \mathcal{C} \neq \emptyset$ and $\alpha \in C$. Assume that $-K_X$ is base point free. Then

$$(5.6) \quad e(\mathcal{M}_H^\alpha(\gamma)^{ss}) = e(\mathcal{M}_H^\mathcal{C}(\gamma)^{ss}) + \sum_{(\gamma_1, \gamma_2, \dots, \gamma_s)} t^{-\sum_{i < j} (c_1(\gamma_i), c_1(\gamma_j))} \prod_{i=1}^s e(\mathcal{M}_H^\mathcal{C}(\gamma_i)^{ss}),$$

where $(\gamma_1, \gamma_2, \dots, \gamma_s)$ satisfy $\mu_\alpha(\gamma_i) = \mu_\alpha(\gamma_j)$ and $\mu_{\alpha'}(\gamma_i) > \mu_{\alpha'}(\gamma_j)$ for $i < j$ and $\alpha' \in C \cap \mathcal{C}$.

Proof. For the Harder-Narasimhan filtration (5.5), Lemma 5.2 implies that

$$(5.7) \quad \text{Ext}^2(F_i/F_{i-1}, F_j/F_{j-1}) = \text{Hom}(F_j/F_{j-1}, F_i/F_{i-1}(K_X))^\vee = 0$$

for $j \leq i$. Moreover we get

$$(5.8) \quad \chi(F_i/F_{i-1}, F_j/F_{j-1}) = \chi(F_j/F_{j-1}, F_i/F_{i-1}) = -(c_1(F_i/F_{i-1}), c_1(F_j/F_{j-1})).$$

Therefore in the same way as in the proof of [Y1, Thm. 3.2] (see the description of the stack of filtration in [Y7, sect. 5.2]), we get this proposition. $\square$

By induction on $(H, c_1(\gamma))$ and Proposition 5.4, we can show the following proposition.

Proposition 5.5. *Assume that $-K_X$ is base point free or K_X is numerically trivial. Then virtual Hodge polynomial $e(\mathcal{M}_H^\alpha(\gamma)^{ss})$ does not depend on the choice of general α .*

As in [E-G], we can construct moduli of α -twisted semi-stable sheaves (see Appendix Theorem 7.1). Hence we get the following theorem.

Theorem 5.6. (i) *There is a coarse moduli scheme $\overline{M}_H^\alpha(\gamma)$ of S -equivalence classes of α -twisted semi-stable sheaves.*
(ii) *$\overline{M}_H^\alpha(\gamma)$ is projective.*
(iii) *For different α, α' , the relation between $\overline{M}_H^\alpha(\gamma)$ and $\overline{M}_H^{\alpha'}(\gamma)$ is described as Mumford-Thaddeus type flips:*

$$(5.9) \quad \begin{array}{ccccccc} \overline{M}_H^{\alpha_1}(\gamma) & & \overline{M}_H^{\alpha_2}(\gamma) & & \overline{M}_H^{\alpha_n}(\gamma) \\ & \searrow & \swarrow & \searrow & \swarrow & \dots & \swarrow \\ & \overline{M}_H^{\alpha_{1,2}}(\gamma) & & \overline{M}_H^{\alpha_{2,3}}(\gamma) & & & \end{array}$$

where $\alpha = \alpha_1, \alpha' = \alpha_n$.

6. FOURIER-MUKAI TRANSFORM ON ELLIPTIC SURFACES

Let $\pi : X \rightarrow C$ be an elliptic surface with a 0-section σ such that every fiber is irreducible. Let f be a fiber of π . Then a compactification of the relative Jacobian of $\pi : X \rightarrow C$ is isomorphic to $\pi : X \rightarrow C$. Let $\mathcal{P}$ be a universal family on $X \times_C X$. Then $\widehat{\mathcal{F}}_{\mathcal{P}}[2]$ is the inverse of $\mathcal{F}_{\mathcal{P}}$.

Let τ be a section of π . We set

$$(6.1) \quad \langle \sigma, f \rangle^\perp := \{D \in \text{NS}(X) \mid (D, \sigma) = (D, f) = 0\}.$$

We shall normalize $\mathcal{P}$ so that $\mathcal{F}_{\mathcal{P}}^1(\mathcal{O}_X) = \mathcal{O}_\sigma$. In [Y8, 3.2], we showed the following:

$$(6.2) \quad \text{For a coherent sheaf } E \text{ of } (\text{rk}(E), c_1(E), -\text{ch}_2(E)) = (r, lf + D, n), D \in \langle \sigma, f \rangle^\perp$$

$$(\text{rk}(\mathcal{F}_{\mathcal{P}}(E)), c_1(\mathcal{F}_{\mathcal{P}}(E)), \chi(\mathcal{F}_{\mathcal{P}}(E))) = -(0, r\sigma + nf - D, r + l).$$

Let $M(r, d) \rightarrow C$ be the relative moduli space of stable sheaves of rank r and degree d .

Lemma 6.1. *Assume that $(r, d) = 1$. Then $M(r, d) \cong X$.*

Proof. We shall prove our claim by induction on r . By tensoring $\mathcal{O}_X(k\sigma)$, we may assume that $0 < d \leq r$. If $r = d$, then $r = d = 1$. Hence our claim obviously holds. If $d < r$, then by $\widehat{\mathcal{F}}_{\mathcal{P}}$, we get an isomorphism $M(r, d) \rightarrow M(d, -r)$. By induction hypothesis, $M(d, -r) \cong X$. Thus our claim holds. $\square$

By this proof, a universal family $\mathcal{E}$ on $M(r, d) \times_C X$ is obtained by compositions of T_σ and $\mathcal{F}_{\mathcal{P}}$ from $\mathcal{P}$ on $X \times_C X$. We consider Fourier-Mukai transform $\mathcal{F}_{\mathcal{E}} : \mathbf{D}(X) \rightarrow \mathbf{D}(X)$ defined by $\mathcal{E}$:

$$(6.3) \quad \mathcal{F}_{\mathcal{E}}(x) = \mathbf{R}p_{2*}(\mathcal{E} \otimes p_1^*(x)), x \in \mathbf{D}(X)$$

where $p_i : X \times X \rightarrow X$, $i = 1, 2$ are two projections.

Lemma 6.2.

$$(6.4) \quad \mathcal{F}_{\mathcal{E}}(\Delta) = \Delta$$

for $\Delta \in \gamma^{-1}(\langle \sigma, f \rangle^\perp)$.

Proof. Since $T_\sigma(D) = (0, D, (\sigma, D)) = (0, D, 0)$. By (6.2), we get $\mathcal{F}_{\mathcal{P}}(\Delta) = \Delta$. Since $\mathcal{E}$ on $M(r, d) \times_C X$ is obtained by compositions of T_σ and $\widehat{\mathcal{F}}_{\mathcal{P}}$ from $\mathcal{P}$, Lemma 1.1 implies our claim. $\square$

We set $E_0 := \widehat{\mathcal{F}}_{\mathcal{E}}^1(\mathcal{O}_\sigma)$ and $F_0 := \mathcal{F}_{\mathcal{E}}^0(\mathcal{O}_\sigma)$. Then E_0 is a vector bundle of $(\text{rk}(E_0), c_1(E_0)) = (r, -d\sigma + kf)$, $k \in \mathbb{Z}$ and F_0 a vector bundle of $(\text{rk}(F_0), c_1(F_0)) = (r, d'\sigma + k'f)$, $k' \in \mathbb{Z}$. In $K(X)$, we see that

$$(6.5) \quad \begin{cases} \mathcal{F}_{\mathcal{E}}(\mathbb{C}) = (F_0)_{|f}, \\ \mathcal{F}_{\mathcal{E}}((E_0)_{|f}) = -\mathbb{C}, \end{cases}$$

where $\mathbb{C}$ is the structure sheaf of $f \cap \sigma$. We see that

$$(6.6) \quad K(X)/\gamma^{-1}(\langle \sigma, f \rangle^\perp) \subset \frac{1}{r}(\mathbb{Z}E_0 + \mathbb{Z}E_{0|f} + \mathbb{Z}\mathbb{C}).$$

For $x = aE_0 + bE_{0|f} + c\mathbb{C}$, $\mathcal{F}_{\mathcal{E}}(x) = -a\mathcal{O}_\sigma - b\mathbb{C} + cF_{0|f}$. Hence

$$(6.7) \quad -\gamma(\mathcal{F}_{\mathcal{E}}(x)) = (0, a\sigma - crf, b - c\chi(F_{0|f}) + a\chi(\mathcal{O}_\sigma)).$$

Since $(r, d) = 1$, if x belongs to $K(X)$, then $a \in \mathbb{Z}$.

Lemma 6.3. *If $c_1 = \lambda f$, $\lambda \in \mathbb{Z}$, then $\mathcal{M}_{\sigma+nf}^\alpha(0, c_1, \chi)^{ss} = \mathcal{M}_{\sigma+nf}(0, c_1, \chi)^{ss}$ for any $\mathbb{Q}$ -divisor α . In particular, $\mathcal{M}_{\sigma+nf}^{F_0}(0, rf, d')^{ss} = \mathcal{M}_{\sigma+nf}(0, rf, d')^{ss}$.*

6.1. Preservation of stability. The following is an easy consequence of the proof of base change theorem (cf. [Y8, Lem. 3.6]).

Lemma 6.4. *Let L be a coherent sheaf of pure dimension 1 on X with $c_1(L) = r\sigma + nf + D$, $D \in \langle \sigma, f \rangle^\perp$. Then L satisfies WIT_1 and $\widehat{\mathcal{F}}_\mathcal{E}^1(L)$ is torsion free, if the following two conditions are satisfied:*

- (1) $\text{Hom}(\mathcal{E}_x, L) = 0$, $x \in X$ except finite subset S of X .
- (2) $\text{Ext}^2(\mathcal{E}_x, L) \cong \text{Hom}(L, \mathcal{E}_x)^\vee = 0$ for all $x \in X$.

Lemma 6.5. *For a purely 1-dimensional sheaf L of $c_1(L) = r\sigma + nf + D$, $D \in \langle \sigma, f \rangle^\perp$, $\text{Hom}_{p_1}(\mathcal{E}, p_2^*L) = 0$.*

Proof. It is sufficient to prove that $\text{Hom}(\mathcal{E}_x, L) = 0$ for some point $x \in X$. We choose a point $x \in X$ which is not contained in $\text{Supp}(L)$. Since L is of pure dimension 1, we get $\text{Hom}(\mathcal{E}_x, L) = 0$. $\square$

Proposition 6.6. *Let L be a coherent sheaf of pure dimension 1 on X with $c_1(L) = x\sigma + nf + D$, $D \in \langle \sigma, f \rangle^\perp$ and $\chi_{F_0}(L) > 0$. If L is F_0 -twisted semi-stable with respect to $\sigma + kf$, $k \gg 0$, then L satisfies two conditions in Lemma 6.4.*

Proof. By taking account of a Jordan-Hölder filtration, we may assume that L is stable. We note that

$$(6.8) \quad \mu_{F_0}(L) := \frac{\chi_{F_0}(L)}{(c_1(L \otimes F_0^\vee), \sigma + kf)} = \frac{\chi_{F_0}(L)}{r((k + (\sigma^2))x + n)}.$$

Since $\mu_{F_0}(\mathcal{E}_x) = 0$, $x \in X$, by Lemma 6.3, (2) holds. So we shall prove (1). Let $D = D_{vir} + D_{hol}$ be the decomposition of the scheme-theoretic support of L , where D_{vir} consists of all fiber components and D_{hol} consists of the other components. Then we have an exact sequence

$$(6.9) \quad 0 \rightarrow F \rightarrow L \rightarrow (L|_{D_{hol}})/T \rightarrow 0,$$

where T is the torsion submodule of $L|_{D_{hol}}$. Then F is a pure dimension 1 subsheaf of L with $c_1(F) = lf$. By the stability of L , we get

$$(6.10) \quad \mu_{F_0}(F) = \frac{\chi_{F_0}(F)}{rl} \leq \frac{\chi_{F_0}(L)}{r((k + (\sigma^2))x + n)}.$$

Since k is sufficiently large (the condition $k > \max\{((c_1(L), \sigma + k_0f)\chi_{F_0}(L) - n)/x - (\sigma^2), k_0\}$ is sufficient, where $\sigma + k_0f$ is ample), we get $\chi_{F_0}(F) \leq 0$. Since $\text{Hom}(\mathcal{E}_x, L|_{D_{hol}}/T) = 0$ for all $x \in X$, we shall prove that $\text{Hom}(\mathcal{E}_x, F) = 0$ except finite numbers of points.

Proof of the claim: Let

$$(6.11) \quad 0 \subset F_1 \subset F_2 \subset \cdots \subset F_s = F$$

be the Harder-Narasimhan filtration of F with respect to $\sigma + kf$. Then

$$(6.12) \quad \mu_{F_0}(F_1) > \mu_{F_0}(F_2/F_1) > \cdots > \mu_{F_0}(F_s/F_{s-1}).$$

Since F_1 is a subsheaf of L , we also have the inequality $\chi_{F_0}(F_1) \leq 0$. If $\text{Hom}(\mathcal{E}_x, F_1) \neq 0$, then $\mu_{F_0}(F_1) = 0$ and F_1 is S -equivalent to $\mathcal{E}_x \oplus E$ for some E . Hence the choice of x is finite. Clearly $\text{Hom}(\mathcal{E}_x, F_i/F_{i-1}) = 0$ for $i \geq 2$. Hence the claim holds. $\square$

Lemma 6.7. *Let E be a torsion free sheaf of $\text{rk}(E) = xr > 0$ and $c_1(E \otimes E_0^\vee) = lf + D$, $D \in \langle \sigma, f \rangle^\perp$ on X . Assume that E is μ -semi-stable with respect to $\sigma + kf$, $k \gg 0$. Then E satisfies WIT_1 and $\mathcal{F}_\mathcal{E}^1(E)$ is of pure dimension 1.*

Proof. We shall first prove that $\mathcal{E} \otimes p_1^*(E)$ is p_2 -flat. Let

$$(6.13) \quad 0 \rightarrow W_1 \rightarrow W_0 \rightarrow \mathcal{E} \rightarrow 0$$

be a locally free resolution of $\mathcal{E}$ on $X \times X$. It is sufficient to prove that

$$(6.14) \quad \psi_x : (W_1)_{|x \times X} \otimes E \rightarrow (W_0)_{|x \times X} \otimes E$$

is injective for all $x \in X$. We note that $\text{rk } W_1 = \text{rk } W_0$ and $\mathcal{E}_x \otimes E$ is a torsion sheaf on X . Since E is torsion free, ψ_x is injective for all $x \in X$. Thus $\mathcal{E} \otimes p_1^*(E)$ is a p_2 -flat sheaf.

Hence we can use base change theorem. Since $p_2 : X \times_C X \rightarrow X$ is relative dimension 1, $R^2 p_{2*}(\mathcal{E} \otimes p_1^*(E)) = 0$. Since $E|_{\pi^{-1}(y)}$ is semi-stable for general $y \in C$, $H^0(X, \mathcal{E}_x \otimes E) = 0$ for a general point x of X . Thus $p_{2*}(\mathcal{E} \otimes p_1^*(E))$ is a torsion sheaf. By the proof of base change theorem, locally there is a complex of locally free sheaves $V_1 \xrightarrow{\phi} V_0$ which is quasi-isomorphic to $\mathbf{R}p_{2*}(\mathcal{E} \otimes p_1^*(E))$. Hence $p_{2*}(\mathcal{E} \otimes p_1^*(E)) = \ker \phi = 0$, which means that E satisfies WIT_1 . Also we get $\text{proj-dim}(\text{coker } \phi) = 1$. Hence $R^1 p_{2*}(\mathcal{E} \otimes p_1^*(E))$ is of pure dimension 1. $\square$

Corollary 6.8. *Let E be a torsion free sheaf on X and assume that $E|_{\pi^{-1}(y)}$ is a semi-stable vector bundle of $\deg(E(-(d/r)\sigma)|_{\pi^{-1}(y)}) = 0$ for a general $y \in C$. Then E satisfies WIT_1 and $\mathcal{F}_\varepsilon^1(E)$ is of pure dimension 1.*

Proof. We consider the Harder-Narasimhan filtration of E with respect to $\sigma + kf$, $k \gg 0$. Applying $\mathcal{F}_\varepsilon$ to this filtration, we get our corollary by Lemma 6.7. $\square$

Lemma 6.9. *We set $\gamma(E) = \gamma(xE_0 + yE_{0|f} + z\mathbb{C}) + D$, $D \in \langle \sigma, f \rangle^\perp$, and assume that E is E_0 -twisted semi-stable with respect to $\sigma + kf$, $k \gg 0$. If $\chi_{F_0}(\mathcal{F}_\varepsilon^1(E)) = x(r\chi(\mathcal{O}_\sigma) - d'(\sigma^2) - k') + ry > 0$, then $\mathcal{F}_\varepsilon^1(E)$ is F_0 -twisted semi-stable.*

Proof. Assume that $\mathcal{F}_\varepsilon^1(E)$ is not semi-stable. Then, there is a stable subsheaf F of $\mathcal{F}_\varepsilon^1(E)$ such that $\mu_{F_0}(F) > \mu_{F_0}(\mathcal{F}_\varepsilon^1(E)) = (\chi_{F_0}(\mathcal{F}_\varepsilon^1(E)))/((k + (\sigma^2))x - rz) > 0$. We set $G := \mathcal{F}_\varepsilon^1(E)/F$. Applying $\widehat{\mathcal{F}}_\varepsilon$ to the exact sequence

$$(6.15) \quad 0 \rightarrow F \rightarrow \mathcal{F}_\varepsilon^1(E) \rightarrow G \rightarrow 0,$$

we get an exact sequence

$$(6.16) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \widehat{\mathcal{F}}_\varepsilon^0(F) & \longrightarrow & 0 & \longrightarrow & \widehat{\mathcal{F}}_\varepsilon^0(G) \\ & & \longrightarrow & \widehat{\mathcal{F}}_\varepsilon^1(F) & \longrightarrow & E & \longrightarrow \widehat{\mathcal{F}}_\varepsilon^1(G) \\ & & & \longrightarrow & \widehat{\mathcal{F}}_\varepsilon^2(F) & \longrightarrow & 0 \longrightarrow \widehat{\mathcal{F}}_\varepsilon^2(G) \longrightarrow 0 \end{array}$$

By Lemma 6.5, $\widehat{\mathcal{F}}_\varepsilon^0(G) = 0$. Since $\mu_{F_0}(F) > 0$, we also get $\widehat{\mathcal{F}}_\varepsilon^2(F) = 0$. Thus F and G satisfies WIT_1 and $\widehat{\mathcal{F}}_\varepsilon^1(F)$ is a subsheaf of E . We set $\gamma(\widehat{\mathcal{F}}_\varepsilon^1(F)) = \gamma(x'E_0 + y'E_{0|f} + z'\mathbb{C}) + D'$, $D' \in \langle \sigma, f \rangle^\perp$. Then $\gamma(F) = (0, x'\sigma - z'r f - D', y' - z'\chi(F_{0|f}) + x'\chi(\mathcal{O}_\sigma))$. Since E is semi-stable with respect to $\sigma + kf$, $k \gg 0$,

- ($\star$) (i) $y'/x' < y/x$, or
- (ii) $y'/x' = y/x$ and $z'/x' \leq z/x$.

On the other hand,

$$(6.17) \quad \begin{aligned} \mu_{F_0}(\mathcal{F}_\varepsilon^1(E)) - \mu_{F_0}(F) &= \frac{xl + yr}{r(k + (\sigma^2))x - rz} - \frac{x'l + y'r}{r(k + (\sigma^2))x' - rz'} \\ &= \frac{(yx' - y'x)r(k + (\sigma^2)) + (x'l + ry)z - (xl + ry)z'}{(r(k + (\sigma^2))x - rz)((k + (\sigma^2))x' - z')} \\ &= \frac{(yx' - y'x)(r(k + (\sigma^2)) - rz/x) + (xl + ry)(x'z/x - z')}{(r(k + (\sigma^2))x - rz)((k + (\sigma^2))x' - z')}, \end{aligned}$$

where $l = r\chi(\mathcal{O}_\sigma) - d'(\sigma^2) - k'$. We note that the choice of x' is finite. In Lemma 6.10, we shall show that the choice of z' is also finite. Then there is an integer $N(x, D, y, z)$ such that for $k > N(x, D, y, z)$,

- (i) $yx' - y'x > 0$ implies $(yx' - y'x)(r(k + (\sigma^2)) - rz/x) + (xl + ry)(x'z/x - z') > 0$ and
- (ii) $yx' - y'x < 0$ implies $(yx' - y'x)(r(k + (\sigma^2)) - rz/x) + (xl + ry)(x'z/x - z') < 0$.

Then ($\star$) implies that $\mu_{F_0}(\mathcal{F}_\varepsilon^1(E)) - \mu_{F_0}(F) \geq 0$, which is a contradiction. Therefore $\mathcal{F}_\varepsilon^1(E)$ is a semi-stable sheaf. $\square$

Lemma 6.10. *Keep the notations as above. Then the choice of z' is finite and the number of such z' is bounded in terms of (x, D, y, z) .*

Proof. We fix an ample divisor $\sigma + k_0f$. Since F is a subsheaf of $\mathcal{F}_\varepsilon^1(E)$,

$$(6.18) \quad 0 \leq (c_1(F), \sigma + k_0f) \leq (c_1(\mathcal{F}_\varepsilon^1(E)), \sigma + k_0f).$$

Since $(c_1(F), \sigma + k_0f) = (k_0 + (\sigma^2))x' - rz'$ and $x' \leq x$, we get our claim. $\square$

Lemma 6.11. *Let L be a pure dimension 1 sheaf of $\gamma(L) = (0, x\sigma - rzf, y - z\chi(F_{0|f}) + x\chi(\mathcal{O}_\sigma))$. Assume that $\chi_{F_0}(L) > 0$ and L is F_0 -twisted semi-stable with respect to $\sigma + kf$, $k \gg 0$. Then $\widehat{\mathcal{F}}_\varepsilon^1(L)$ is an E_0 -twisted semi-stable sheaf with respect to $\sigma + kf$, $k \gg 0$.*

Proof. We note that Lemma 6.4 and Proposition 6.6 imply that L satisfies WIT_1 and $\widehat{\mathcal{F}}_\varepsilon^1(L)$ is torsion free. Assume that $\widehat{\mathcal{F}}_\varepsilon^1(L)$ is not semi-stable with respect to $\sigma + kf$, $k \gg 0$. Then there is a destabilizing subsheaf F of $\widehat{\mathcal{F}}_\varepsilon^1(L)$. We set $G := \widehat{\mathcal{F}}_\varepsilon^1(L)/F$. It is easy to see that $\widehat{\mathcal{F}}_\varepsilon^1(L)|_{\pi^{-1}(y)}$ is semi-stable for general $y \in C$. Since k is sufficiently large, $F|_{\pi^{-1}(y)}$ and $G|_{\pi^{-1}(y)}$ are semi-stable vector bundles of degree 0 for general $y \in C$. Then Corollary 6.8 implies that F and G satisfy WIT_1 and we get an exact sequence

$$(6.19) \quad 0 \rightarrow \mathcal{F}_\varepsilon^1(F) \rightarrow L \rightarrow \mathcal{F}_\varepsilon^1(G) \rightarrow 0.$$

In the same way as in Lemma 6.9, we get a contradiction. Thus $\widehat{\mathcal{F}}_\varepsilon^1(L)$ is semi-stable. $\square$

Therefore, we get the following theorem.

Theorem 6.12. *Keep the notation as above. For $x \in K(X)$ of $\text{rk } x > 0$, Assume that the relative twisted degree $(c_1(x \otimes E_0^\vee), f) = 0$. Then $\mathcal{F}_\mathcal{E}$ gives an isomorphism of moduli stack*

$$(6.20) \quad \mathcal{M}_{\sigma+kf}^{E_0}(\gamma(x))^{ss} \rightarrow \mathcal{M}_{\sigma+kf}^{F_0}(-\gamma(\mathcal{F}_\mathcal{E}(x)))^{ss}$$

if $\chi_{F_0}(-\mathcal{F}_\mathcal{E}(x)) > 0$ and $k \gg 0$.

Remark 6.1. Let $\varepsilon \in \langle \sigma, f \rangle^\perp$ be a $\mathbb{Q}$ -divisor of $-(\varepsilon^2) \ll 1$. Then we can show that

$$(6.21) \quad \mathcal{M}_{\sigma+kf+\varepsilon}^{E_0}(\gamma(x))^{ss} \rightarrow \mathcal{M}_{\sigma+kf}^{F_0(-\varepsilon)}(-\gamma(\mathcal{F}_\mathcal{E}(x)))^{ss}$$

if $(c_1(x \otimes E_0^\vee), f) = 0$, $\chi_{F_0}(-\mathcal{F}_\mathcal{E}(x)) > 0$ and $k \gg 0$.

6.2. Application of Theorem 6.12.

Lemma 6.13. *Let $H = \sigma + kf$ be an ample divisor. We set $\gamma_n := (0, \sigma + nf + D, \chi)$, $D \in \langle \sigma, f \rangle$. Then for a general $t \in \mathbb{Q}$, $\mathcal{M}_H^{t(\sigma+mf)}(\gamma_n)^{ss} = \mathcal{M}_H^{t(\sigma+mf)}(\gamma_n)^s$, if $m \neq k$.*

Proposition 6.14. *We set $\gamma_n := (0, \sigma + nf + D, \chi)$, $D \in \langle \sigma, f \rangle^\perp$. Then $e(M_H^\alpha(\gamma_n))$ does not depend on α , if α does not lie on walls.*

Proof. It is sufficient to prove the claim for $\mathcal{M}_H^\alpha(\gamma_n)^{ss}$. We prove our claim by induction on n . We note that there is a section τ of π such that $D = \tau - \sigma - (\tau - \sigma, \sigma)f$. Hence $\sigma + nf + D = \tau + n'f$, $n' = n - (\tau - \sigma, \sigma)$. Let W be a wall and α belongs to W . Let α_+, α_- be $\mathbb{Q}$ -divisors which are very close to α and $\alpha = (\alpha_+ + \alpha_-)/2$. We consider Harder-Narasimhan filtration (5.5) in Proposition 5.1, where $\alpha_1 = \alpha_+, \alpha_-$. Since $c_1(E) = \tau + n'f$, we get $c_1(F_i/F_{i-1}) = l_i f, \tau + l_i f$. Assume that $c_1(F_i/F_{i-1}) = l_i f$ and $c_1(F_j/F_{j-1}) = l_j f$ for different i and j . Since $\mu_\alpha(F_i/F_{i-1}) = \mu_\alpha(F_j/F_{j-1})$, $(F_i/F_{i-1})/l_i = (F_j/F_{j-1})/l_j$ in $K(X) \otimes \mathbb{Q}$. Then $\mu_{\alpha_\pm}(F_i/F_{i-1}) = \mu_{\alpha_\pm}(F_j/F_{j-1})$, which is a contradiction. Therefore $s = 2$ and $c_1(F_i/F_{i-1}) = l_i f$ and $c_1(F_j/F_{j-1}) = \tau + l_j f$, $\{i, j\} = \{1, 2\}$. Since F_1 or F_2/F_1 is supported on some fibers, we see that

$$(6.22) \quad \text{Ext}^2(F_2/F_1, F_1) = \text{Hom}(F_1, F_2/F_1(K_X))^\vee = \text{Hom}(F_1, F_2/F_1)^\vee = 0.$$

Since $-\chi(F_2/F_1, F_1) = (c_1(F_2/F_1), c_1(F_1))$, by the same argument as in Proposition 5.4, we get

$$(6.23) \quad e(\mathcal{M}_H^\alpha(\gamma_n)^{ss}) = e(\mathcal{M}_H^{\alpha_\pm}(\gamma_n)^{ss}) + \sum_k e(\mathcal{M}_H^{\alpha_\pm}(0, \tau + (n' - kl)f, \chi - kd)^{ss})e(\mathcal{M}_H^{\alpha_\pm}(0, kl f, kd)^{ss})t^{kl}$$

where (l, d) satisfies that $\mu_\alpha(\gamma_n) = (d - l(\alpha, f))/l(H, f)$. By induction hypothesis,

$$(6.24) \quad e(\mathcal{M}_H^{\alpha_+}(0, \tau + (n' - kl)f, \chi - kd)^{ss}) = e(\mathcal{M}_H^{\alpha_-}(0, \tau + (n' - kl)f, \chi - kd)^{ss}).$$

Also we know that $\mathcal{M}_H^\beta(0, kl f, kd)^{ss}$ does not depend on β . Therefore we get our claim. $\square$

The following is a generalization of [Gö] and [Y2].

Theorem 6.15. *If $(r, (c_1, f)) = 1$, then*

$$(6.25) \quad e(M_{\sigma+kf}(r, c_1, \chi)) = e(\text{Pic}^0(X) \times \text{Hilb}_X^n),$$

where $2n + h^1(\mathcal{O}_X) = \dim M_{\sigma+kf}(r, c_1, \chi)$ and $k \gg 0$.

Proof. Let x be an element of $K(X)$ such that $\gamma(x) = (r, c_1, \chi)$. We set $d := (c_1, f)$. We consider Fourier-Mukai transform induced by a universal family $\mathcal{E}$ on $X \times_C M(r, d) = X \times_C X$. Then $-c_1(\mathcal{F}_\mathcal{E}(x)) = \sigma + nf + D$, $n \in \mathbb{Z}$, $D \in \langle \sigma, f \rangle^\perp$. Replacing x by $x \otimes \mathcal{O}_X(mf)$, $m \gg 0$, we may assume that $\chi(-\mathcal{F}_\mathcal{E}(x)) > 0$ and $\chi_{F_0}(-\mathcal{F}_\mathcal{E}(x)) > 0$. By Theorem 6.12, $M_{\sigma+kf}(\gamma(x))$ is isomorphic to $M_{\sigma+kf}^\alpha(-\gamma(\mathcal{F}_\mathcal{E}(x)))$. By Proposition 6.14, $e(M_{\sigma+kf}^\alpha(-\gamma(\mathcal{F}_\mathcal{E}(x)))) = e(M_{\sigma+kf}^0(-\gamma(\mathcal{F}_\mathcal{E}(x))))$. By using Theorem 6.12 again, we see that $M_{\sigma+kf}^0(-\gamma(\mathcal{F}_\mathcal{E}(x)))$ is isomorphic to Hilbert scheme of points. $\square$

By the similar arguments as in the proof of Proposition 6.14, we also get the following proposition.

Proposition 6.16. *$e(M_H^\alpha(\gamma_n))$ does not depend on the choice of H , if H is general.*

Proposition 6.17. *We set $\gamma_n = (0, \tau + nf, m)$, where τ is a section of π . Then $M_H^\alpha(\gamma_n)$ is smooth for general α and H .*

Proof. Let F be a simple pure dimension 1 sheaf of $c_1(F) = \tau + nf$. It is sufficient to show that $H^0(X, K_X) \rightarrow \text{Hom}(F, F(K_X))$ is an isomorphism. We set $F_2 := F|_\sigma / (\text{torsion})$ and $F_1 := \ker(F \rightarrow F_2)$. Since F_1 is supported on fibers, $F_1(K_X) \cong F_1$. By the simpleness of F , $\text{Hom}(F, F_1(K_X)) \cong \text{Hom}(F, F_1) = 0$. We note that $\text{Hom}(F_1, F_2(K_X)) = 0$. Hence $\text{Hom}(F, F(K_X)) \rightarrow \text{Hom}(F, F_2(K_X))$ is injective and $\text{Hom}(F_2, F_2(K_X)) \rightarrow \text{Hom}(F, F_2(K_X))$ is isomorphic. Since F_2 is a line bundle on τ ,

$$(6.26) \quad H^0(C, \pi_*(K_X)) = H^0(X, K_X) \rightarrow \text{Hom}(F_2, F_2(K_X))$$

is isomorphic. Therefore $H^0(X, K_X) \rightarrow \text{Hom}(F, F(K_X))$ must be isomorphic. $\square$

From now on, we assume that X is a rational elliptic surface. By using Proposition 5.5 and Remark 6.1, we also get the following.

Corollary 6.18. *Assume that X is a rational elliptic surface. Let $\varepsilon, \varepsilon' \in \langle \sigma, f \rangle^\perp$ be general $\mathbb{Q}$ -divisors of $-(\varepsilon^2), -(\varepsilon'^2) \ll 1$. If $\gcd(r, (c_1, f)) = l$, then*

$$(6.27) \quad e(\mathcal{M}_{\sigma+kf+\varepsilon}(r, c_1, \chi)^{ss}) = e(\mathcal{M}_{\sigma+kf+\varepsilon'}(l, \xi, \chi')^{ss}), k \gg 0$$

for some $\chi' \in \mathbb{Z}$ and $\xi \in H^2(X, \mathbb{Z})$ of $(\xi, f) = 0$. In particular, if $l = 2$, then we obtain $e(\mathcal{M}_{\sigma+kf+\varepsilon}(r, c_1, \chi)^{ss})$ from [Y5].

We also get the following corollary which is a generalization of [Y5].

Corollary 6.19. *Keep notation as in Corollary 6.18. If (r, D, χ) , $D \in \langle \sigma, f \rangle^\perp$ is primitive, then*

$$(6.28) \quad e(\mathcal{M}_{\sigma+kf+\varepsilon}(r, D, \chi)^{ss}) = e(\mathcal{M}_{\sigma+kf+\varepsilon'}(r, D + tf, \chi)^{ss}), k \gg 0$$

for $t \in \mathbb{Z}$.

Proof. If (r, D, χ) , $D \in \langle \sigma, f \rangle^\perp$ is primitive, then $C := r\sigma + (r - \chi)f - D$ is primitive. Since $\text{NS}(X) = H^2(X, \mathbb{Z})$ is a unimodular lattice, there is a divisor β such that $(C, \beta) = 1$. Hence $e(\mathcal{M}_{\sigma+kf}^{\alpha+t\beta}(0, C, m)^{ss}) = e(\mathcal{M}_{\sigma+kf}^{\alpha}(0, C, m - t)^{ss})$, $t \in \mathbb{Z}$. Therefore $e(\mathcal{M}_{\sigma+kf+\varepsilon}(r, D, \chi)^{ss}) = e(\mathcal{M}_{\sigma+kf+\varepsilon'}(r, D + tf, \chi)^{ss})$. $\square$

7. APPENDIX

Let X be a projective surface defined over a field k . Let α be a $\mathbb{Q}$ -divisor on X . In this appendix, we shall prove the following.

Theorem 7.1. *There is a projective moduli scheme $\overline{M}_H^\alpha(\gamma)$ of α -twisted semi-stable sheaves of pure dimension 1 on X with respect to H .*

Since α -twisted semi-stability does not change under the operation $E \mapsto E \otimes \mathcal{O}_X(nH)$, we may assume that $\alpha = D/n$ for some positive integer n and an irreducible divisor D on X of $(D, H) > (c_1, H)$. We note that

$$(7.1) \quad \begin{aligned} \chi(E(-\alpha)) &= \frac{1}{n} \chi(E(-D)) + \left(1 - \frac{1}{n}\right) \chi(E) \\ &= \chi(E(-D)) + \left(1 - \frac{1}{n}\right) \chi(E|_D). \end{aligned}$$

As in [E-G], we can regard $\chi(E(-\alpha))$ as the parabolic Euler characteristic of the parabolic sheaf $E(-D) \subset E$, where the weight is $1 - 1/n$. Then it is sufficient to construct moduli space of parabolic semi-stable sheaves on X . The moduli space of parabolic stable sheaves was constructed by Maruyama and Yokogawa [Ma-Yk] and Inaba [I]. They constructed moduli space as GIT quotient of a suitable space. Then the problem is to analyse properly semi-stable points of this space. For a nonsingular variety, Yokogawa [Yk] constructed the moduli space of parabolic semi-stable torsion free sheaves. By technical reason, we can only treat the dimension 1 case.

7.1. Construction of moduli spaces. Let $(X, \mathcal{O}_X(1))$ be a projective scheme X over a field k and an ample line bundle $\mathcal{O}_X(1)$ on X . Let us start with the definition of parabolic sheaves.

Definition 7.1. Let E be a pure dimensional 1 coherent sheaf on X such that $\dim(D \cap \text{Supp}(E)) = 0$. Let

$$(7.2) \quad F(E) : E(-D) = F_{l+1}(E) \subset F_l(E) \subset \cdots \subset F_2(E) \subset F_1(E) = E$$

be a filtration of coherent sheaves and $0 < \alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_l \leq 1$ a sequence of rational numbers. Then the triple $E_* := (E, F(E), \alpha_*)$ is called a parabolic sheaf on X , where $\alpha_* = (\alpha_1, \alpha_2, \dots, \alpha_l)$.

We shall generalize the notion of parabolic sheaves as follows.

Definition 7.2. Let E be a purely 1 dimensional coherent sheaf on X . Let

$$(7.3) \quad F(E) : F_{l+1}(E) \subset F_l(E) \subset \cdots \subset F_2(E) \subset F_1(E) = E$$

be a filtration of coherent sheaves such that $\dim(E/F_{l+1}(E)) = 0$ and $0 < \alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_l \leq 1$ a sequence of rational numbers. Then we call the triple $E_* := (E, F(E), \alpha_*)$ a generalized parabolic sheaf on X .

Thus a generalized parabolic sheaf is a sheaf with a special type of filtration and a sequence of rational numbers.

Definition 7.3. Let $E_* := (E, F(E), \alpha_*)$ be a generalized parabolic sheaf. We set

$$(7.4) \quad \text{par-}\chi(E_*(m)) := \chi(F_{l+1}(m)) + \sum_{i=1}^l \alpha_i \chi(gr_i(E)(m)),$$

where $gr_i(E) = F_i(E)/F_{i+1}(E)$.

We set $\varepsilon_i := \alpha_{i+1} - \alpha_i$, $1 \leq i \leq l$, where $\alpha_{l+1} = 1$. Then

$$(7.5) \quad \text{par-}\chi(E_*(m)) = \chi(E(m)) - \sum_{i=1}^l \varepsilon_i \chi(gr_i(E)(m)).$$

For a coherent sheaf E of dimension 1, we define $a_1(E) \in \mathbb{Z}$ by

$$(7.6) \quad \chi(E(m)) = a_1(E)m + \chi(E).$$

For a numerical polynomial of degree 1

$$(7.7) \quad h(x) = a_1 x + a_0, a_i \in \mathbb{Z},$$

we set $a_i(h) := a_i$.

Definition 7.4. A generalized parabolic sheaf E_* of dimension 1 is sem-stable (resp. stable) with respect to $\mathcal{O}_X(1)$, if

$$(7.8) \quad \frac{\text{par-}\chi(E'_*)}{a_1(E')} \leq \frac{\text{par-}\chi(E_*)}{a_1(E)} \quad (<)$$

for all non-trivial generalized parabolic subsheaf E'_* of E_* .

As in [Ma-Yk], we can consider Jordan-Hölder filtration and define S -equivalence classes of generalized parabolic semi-stable sheaves.

Let $h(x)$ be a numerical polynomial of degree 1 and $h_i(x)$, $1 \leq i \leq l$ constant numerical polynomials. We set $h_*(x) := (h(x), h_1(x), \dots, h_l(x))$. We shall construct the moduli space $\overline{M}_{X/\mathbb{C}}^{h_*, \alpha_*}$ of S -equivalence classes of generalized parabolic semi-stable sheaves $(E, F(E), \alpha_*)$ of Hilbert polynomials

$$(7.9) \quad (\chi(E(m)), \chi(gr_1(E)(m)), \dots, \chi(gr_l(E)(m))) = h_*(m).$$

Let D be a Cartier divisor on X . Then we can construct the moduli space $\overline{M}_{D/X/\mathbb{C}}^{h_*, \alpha_*}$ of S -equivalence classes of parabolic semi-stable sheaves as a locally closed subscheme of $\overline{M}_{X/\mathbb{C}}^{h_*, \alpha_*}$.

Definition 7.5. Let λ be a rational number. Then a purely 1-dimensional sheaf E is of type λ , if

$$(7.10) \quad \frac{\chi(E'')}{a_1(E'')} \geq \frac{\chi(E)}{a_1(E)} - \lambda$$

for all quotient sheaf E'' of pure dimension 1.

Lemma 7.2. Let $E_* = (E, F(E), \alpha_*)$ be a generalized parabolic semi-stable sheaf. Then E is of type $\sum_i \varepsilon_i \chi(gr_i(E))/a_1(E)$.

Proof. Let $E \rightarrow E''$ be a quotient such that E'' is of pure dimension 1. Let E''_* be the induced generalized parabolic structure. Then

$$(7.11) \quad \frac{\chi(E'') - \sum_i \varepsilon_i \chi(gr_i(E''))}{a_1(E'')} \geq \frac{\chi(E) - \sum_i \varepsilon_i \chi(gr_i(E))}{a_1(E)}.$$

Hence $\chi(E'')/a_1(E'') \geq \chi(E)/a_1(E) - \sum_i \varepsilon_i \chi(gr_i(E))/a_1(E)$. $\square$

Let E_* be a generalized parabolic sheaf such that E is of type λ . Let $E_* \rightarrow E''_*$ be a quotient generalized parabolic sheaf such that E'' is of pure dimension 1 and

$$(7.12) \quad \frac{\chi(E) - \sum_i \varepsilon_i \chi(\text{gr}_i(E))}{a_1(E)} \geq \frac{\chi(E'') - \sum_i \varepsilon_i \chi(\text{gr}_i(E''))}{a_1(E'')}.$$

Then $\chi(E)/a_1(E) + \sum_i \varepsilon_i \chi(\text{gr}_i(E)) \geq \chi(E'')/a_1(E'')$. Since the set of E_* is bounded, by Grothendieck's boundedness theorem, the set of such quotients E''_* is bounded. Hence there is an integer $m(\lambda)$ which depends on h_*, α_* and λ such that, for $m \geq m(\lambda)$ and the kernel E'_* of $E_* \rightarrow E''_*$ which satisfies (7.12),

- (b1) $F_i(E')(m)$, $1 \leq i \leq l+1$ are generated by global sections and
- (b2) $H^1(X, F_i(E')(m)) = 0$.

In particular,

- (i) $E(m)$ is generated by global sections and $H^1(X, E(m)) = 0$,
- (ii) $H^1(X, F_i(E)(m)) = 0$.

Let V_m be a vector space of dimension $h(m)$. Let $\mathcal{Q} := \text{Quot}_{V_m \otimes \mathcal{O}_X/X}^{h[m]}$ be the quot-scheme and $V_m \otimes \mathcal{O}_{\mathcal{Q} \times X} \rightarrow \tilde{E}$ the universal quotient sheaf, where $h[m](x) = h(m+x)$. We set $\mathcal{Q}_i := \text{Quot}_{\tilde{E}/\mathcal{Q} \times X/\mathcal{Q}}^{h_i[m]}$, $1 \leq i \leq l$ and let $\tilde{E} \otimes \mathcal{O}_{\mathcal{Q}_i \times X} \rightarrow \tilde{E}_i$ be the universal quotient sheaf. Then there is a closed subscheme Γ of $\prod_{i=1}^l \mathcal{Q}_i$ which parametrizes sequences of quotients

$$(7.13) \quad V_m \otimes \mathcal{O}_X \rightarrow E(m) \rightarrow E_l(m) \rightarrow E_{l-1}(m) \rightarrow \cdots \rightarrow E_1(m) \rightarrow 0$$

such that $V_m \otimes \mathcal{O}_X \rightarrow E(m) \in \mathcal{Q}$ and $E(m) \rightarrow E_i(m) \in \mathcal{Q}_i$, $1 \leq i \leq l$. We set $F_{i+1}(E) := \ker(E \rightarrow E_i)$, $1 \leq i \leq l$ and $F_1(E) := E$. Then we have a filtration

$$(7.14) \quad F(E) : F_{l+1}(E) \subset F_l(E) \subset \cdots \subset F_1(E) = E.$$

Thus $F(E)$ and α_* give a structure of generalized parabolic sheaf on E .

Let Γ^{ss} be the open subscheme of Γ consisting of quotients (7.13) such that

- (i) $V_m \rightarrow H^0(X, E(m))$ is an isomorphism and
- (ii) $(E, F(E), \alpha_*)$ is a generalized parabolic semi-stable sheaf.

Let $G(n) := \text{Gr}(V_m \otimes W, h[m](n))$ be the Grassmannian parametrizing $h[m](n)$ -dimensional quotient spaces of $V_m \otimes W$. We set $G_i := \text{Gr}(V_m, H_i(m))$. Assume that $m \geq m(\lambda)$, $\lambda \geq \sum_i \varepsilon_i a_0(h)/a_1(h)$. Then by Lemma 7.2 and (b1, 2), all generalized parabolic semi-stable sheaves E_* are parametrized by Γ^{ss} , $H^0(X, E(m)) \rightarrow H^0(X, E_i(m))$ is surjective and $H^1(X, E_i(m)) = 0$. For a sequence of quotients

$$(7.15) \quad V_m \otimes \mathcal{O}_X \rightarrow E(m) \rightarrow E_l(m) \rightarrow E_{l-1}(m) \rightarrow \cdots \rightarrow E_1(m) \rightarrow 0 \in \Gamma^{ss},$$

quotient vector spaces $\alpha : V_m \otimes W \rightarrow H^0(X, E(m+n))$ and $\alpha_i : V_m \rightarrow H^0(X, E_i(m))$ define a point of $G(n) \times \prod_i G_i$. Thus we get a morphism $\Gamma^{ss} \rightarrow G(n) \times \prod_i G_i$. As in [I], we can show that this morphism is an immersion

$$(7.16) \quad \Gamma^{ss} \hookrightarrow G(n) \times \prod_i G_i.$$

$SL(V_m)$ acts on $G(n) \times \prod_i G_i$. Let $\mathcal{O}_{G(n)}(1)$ and $\mathcal{O}_{G_i}(1)$, $1 \leq i \leq l$ be tautological line bundles on $G(n)$ and G_i respectively. These line bundles have $SL(V_m)$ -linearizations. We consider GIT semi-stability with respect to a $\mathbb{Q}$ -line bundle $L = \mathcal{O}_{G(n)}(\beta_0) \otimes \mathcal{O}_{G_1}(\beta_1) \otimes \cdots \otimes \mathcal{O}_{G_l}(\beta_l)$.

Proposition 7.3. [I, Prop. 3.2] *Let $\alpha : V_m \otimes W \rightarrow A$ and $\alpha_i : V_m \rightarrow A_i$ be quotients corresponding to a point of $G(n) \times \prod_i G_i$. Then it is GIT semi-stable with respect to L if and only if*

$$(7.17) \quad \dim V_m(\beta_0 \dim \alpha(V' \otimes W) + \sum_i \beta_i \dim \alpha_i(V')) - \dim V'(\beta_0 \dim \alpha(V_m \otimes W) + \sum_i \beta_i \dim \alpha_i(V_m)) \geq 0$$

for all non-zero subspaces V' of V_m .

We set $\beta_0 := (h(m) - \sum_{i=1}^l \varepsilon_i h_i(m))/a_1(h)n$, $\beta_i := \varepsilon_i$ for $1 \leq i \leq l$. We also set $V_i := \ker(\alpha_i|_{V'})$. Then

$$\begin{aligned}
& \dim V_m(\beta_0 \dim \alpha(V' \otimes W) + \sum_i \beta_i \dim \alpha_i(V')) - \dim V'(\beta_0 \dim \alpha(V_m \otimes W) + \sum_i \beta_i \dim \alpha_i(V_m)) \\
&= h(m) \left(\frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} \dim \alpha(V' \otimes W) + \sum_i \varepsilon_i (\dim V' - \dim V_i) \right) \\
&\quad - \dim V' \left(\frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} h[m](n) + \sum_i \varepsilon_i h_i(m) \right) \\
(7.18) \quad &= h(m) \left(\frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} \dim \alpha(V' \otimes W) + \sum_i \varepsilon_i (\dim V' - \dim V_i) \right. \\
&\quad \left. - \dim V' - \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n} \right) \\
&= h(m) \left(\frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} \dim \alpha(V' \otimes W) - \sum_i \varepsilon_i \dim V_i - \alpha_1 \dim V' \right. \\
&\quad \left. - \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n} \right).
\end{aligned}$$

Let $(G(n) \times \prod_i G_i)^{ss}$ be the open subscheme of $G(n) \times \prod_i G_i$ consisting of GIT semi-stable points. Then we get the following.

Proposition 7.4. *We set $\beta_0 := (h(m) - \sum_{i=1}^l \varepsilon_i h_i(m))/a_1(h)n$, $\beta_i := \varepsilon_i$ for $1 \leq i \leq l$. Then there is an integer m_1 such that for all $m \geq m_1$, Γ^{ss} is contained in $(G(n) \times \prod_i G_i)^{ss}$, where $n \gg m$.*

Proof. We set

$$(7.19) \quad \mathcal{F} := \{E' \subset E(m) \mid E' \text{ is generated by } V' \subset V\}.$$

Since $\mathcal{F}$ is a bounded set, for a sufficiently large n which depends on m ,

- (i) $\alpha(V' \otimes W) = H^0(X, E'(n))$, $H^i(X, E'(n)) = 0$, $i > 0$ and
- (ii)

$$(7.20) \quad \left| \frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} (\chi(E') - \dim V') \right| < \frac{1}{a_1(h)}.$$

Then $\dim \alpha(V' \otimes W) = \chi(E'(n)) = a_1(E')n + \chi(E')$. Since E_* is semi-stable, in the same way as in [Ma-Yk, Prop. 2.5], we see that there is an integer m_0 such that for $m \geq m_0$ and a generalized parabolic subsheaf E'_* of $E_*(m)$,

$$(7.21) \quad \frac{\alpha_1 h^0(F_{l+1}(E')) + \sum_{i=1}^l \varepsilon_i h^0(F_i(E'))}{a_1(E')} \leq \frac{\alpha_1 h^0(F_{l+1}(E(m))) + \sum_{i=1}^l \varepsilon_i h^0(F_i(E(m)))}{a_1(E)}$$

and the equality holds, if and only if $\text{par-}\chi(E'_*)/a_1(E') = \text{par-}\chi(E_*)/a_1(E)$. Hence if the equality holds, then E'_* is semi-stable and we may assume that (b1, 2) holds for $E'_*(-m)$. In particular, $\dim V' = \chi(E')$.

We note that

$$\begin{aligned}
& \frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} \dim \alpha(V' \otimes W) - \sum_i \varepsilon_i \dim V_i - \alpha_1 \dim V' - \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n} \\
(7.22) \quad &= (h(m) - \sum_i \varepsilon_i h_i(m)) \frac{a_1(E')}{a_1(E)} - (\sum_i \varepsilon_i \dim V_i + \alpha_1 \dim V') + \frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} (\chi(E') - \dim V').
\end{aligned}$$

By (7.18) and (7.22), if the inequality in (7.21) is strict, then we get

$$(7.23) \quad \dim V_m(\beta_0 \dim \alpha(V' \otimes W) + \sum_i \beta_i \dim \alpha_i(V')) - \dim V'(\beta_0 \dim \alpha(V_m \otimes W) + \sum_i \beta_i \dim \alpha_i(V_m)) > 0.$$

If the equality holds in (7.21), then $\dim V' = \chi(E')$, and hence (7.22) implies that L.H.S. of (7.23) is 0. Therefore our claim holds. $\square$

Remark 7.1. If $\beta_0 = (h(m) - \sum_i \varepsilon_i h_i(m))/(a_1(h)n + t)$, $t \in \mathbb{Q}$, then we can show that $\Gamma^{ss} \cap (G(n) \times \prod_i G_i)^{ss}$ parametrizes generalized semi-stable parabolic sheaves E_* such that

$$(7.24) \quad t \frac{\chi(E)}{a_1(E)} \leq t \frac{\chi(E')}{a_1(E')}$$

for all generalized parabolic subsheaves E'_* of $\text{par-}\chi(E'_*)/a_1(E') = \text{par-}\chi(E_*)/a_1(E)$. If $t = h(m)$, then L is nothing but the polarization in [I].

Proposition 7.5. *There is an integer m_2 such that for all $m \geq m_2$, Γ^{ss} is a closed subscheme of $(G(n) \times \prod_i G_i)^{ss}$, where $n \gg m$.*

Proof. We choose an m so that $h(m)/a_1(h) - \sum_i \varepsilon_i h_i(m) > 0$. We shall prove that $\Gamma^{ss} \rightarrow (G(n) \times \prod_i G_i)^{ss}$ is proper. Let $(R, \mathfrak{m})$ be a discrete valuation ring and K the quotient field of R . We set $T := \text{Spec}(R)$ and $U := \text{Spec}(K)$. Let $U \rightarrow \Gamma^{ss}$ be a morphism such that $U \rightarrow \Gamma^{ss} \rightarrow (G(n) \times \prod_i G_i)^{ss}$ is extended to a morphism $T \rightarrow (G(n) \times \prod_i G_i)^{ss}$. Since Γ is a closed subscheme of $\prod_i \mathcal{Q}_i$, there is a morphism $T \rightarrow \Gamma$, i.e., there is a flat family of a sequence of quotients

$$(7.25) \quad V \otimes \mathcal{O}_{T \times X} \rightarrow \mathcal{E}(m) \rightarrow \mathcal{E}_l(m) \rightarrow \mathcal{E}_{l-1}(m) \rightarrow \cdots \rightarrow \mathcal{E}_1(m) \rightarrow 0.$$

Let $\alpha : V_m \otimes W \otimes R \rightarrow p_{T*}(\mathcal{E}(m+n))$ and $\alpha_i : V_m \otimes R \rightarrow A_i$ be the quotient bundles of $V_m \otimes W \otimes R$ and $V_m \otimes R$ corresponding to the morphism $T \rightarrow (G(n) \times \prod_i G_i)^{ss}$. We set $E := \mathcal{E} \otimes R/\mathfrak{m}$, $E_i := \mathcal{E}_i \otimes R/\mathfrak{m}$ and $F_{i+1}(E) := \ker(E \rightarrow E_i)$.

Claim 7.1. $V_m \rightarrow H^0(X, E(m))$ is injective.

Indeed, we set $V' := \ker(V_m \rightarrow H^0(X, E(m)))$. Then $\alpha(V' \otimes W) = 0$. By (7.18),

$$(7.26) \quad \begin{aligned} 0 &\leq \dim V_m (\beta_0 \dim \alpha(V' \otimes W) + \sum_i \beta_i \dim \alpha_i(V')) - \dim V' (\beta_0 \dim \alpha(V_m \otimes W) + \sum_i \beta_i \dim \alpha_i(V_m)) \\ &= h(m) \left(- \sum_i \varepsilon_i \dim V_i - \alpha_1 \dim V' - \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n} \right) \\ &\leq -h(m) \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n}. \end{aligned}$$

Therefore $V' = 0$.

Claim 7.2. There is a rational number λ which depends on h_* and α_* such that E is of type λ .

Proof of the claim: Let $E \rightarrow E''$ be a quotient of E . Let E' be the kernel of $E \rightarrow E''$. We note that $V_m \rightarrow H^0(X, E(m))$ is injective. We set $V' := V_m \cap H^0(X, E'(m))$. Then $h^0(X, E''(m)) \geq \dim V_m - \dim V'$. By (7.18),

$$(7.27) \quad \begin{aligned} 0 &\leq \dim V (\beta_0 \dim \alpha(V' \otimes W) + \sum_i \beta_i \dim \alpha_i(V')) - \dim V' (\beta_0 \dim \alpha(V \otimes W) + \sum_i \beta_i \dim \alpha_i(V)) \\ &= h(m) \left(\frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim \alpha(V' \otimes W)}{a_1(h)n} + \sum_i \varepsilon_i (\dim V' - \dim V_i) - \dim V' \right. \\ &\quad \left. - \frac{(h(m) - \sum_i \varepsilon_i h_i(m)) \dim V'}{a_1(h)n} \right). \end{aligned}$$

Let F be a subsheaf of $E(m)$ generated by V' . Then F belongs to $\mathcal{F}$. Let ε be a positive number such that $h(m)/a_1(h) - \sum_i \varepsilon_i h_i(m) - \varepsilon > 0$. Since $\mathcal{F}$ is a bounded set, for a sufficiently large n which depends on m and ε , we have $\alpha(V' \otimes W) = H^0(X, F(n))$, $H^1(X, F(n)) = 0$ and

$$(7.28) \quad \left| \left(h(m) - \sum_i \varepsilon_i h_i(m) \right) \frac{\dim \alpha(V' \otimes W)}{a_1(h)n} - \left(h(m) - \sum_i \varepsilon_i h_i(m) \right) \frac{a_1(F)}{a_1(h)} \right| < \varepsilon.$$

Therefore

$$(7.29) \quad \begin{aligned} 0 &\leq \frac{h(m) - \sum_i \varepsilon_i h_i(m)}{a_1(h)n} \dim \alpha(V' \otimes W) + \sum_i \varepsilon_i (\dim V' - \dim V_i) - \dim V' \\ &< (h(m) - \sum_i \varepsilon_i h_i(m)) \frac{a_1(F)}{a_1(h)} + \varepsilon + \sum_i \varepsilon_i (\dim V' - \dim V_i) - \dim V', \end{aligned}$$

where $V_i = \ker(\alpha_i|_{V'})$. Since $\dim(V'/V_i) \leq \dim(\text{im } \alpha_i) = h_i(m)$, we get

$$(7.30) \quad h(m) \frac{a_1(F)}{a_1(h)} - \dim V' > \sum_i \varepsilon_i h_i(m) \left(\frac{a_1(F)}{a_1(h)} - 1 \right) - \varepsilon \geq - \sum_i \varepsilon_i h_i(m) - \varepsilon.$$

Since $a_1(E') \geq a_1(F)$,

$$\begin{aligned}
(7.31) \quad \frac{h^0(X, E''(m))}{a_1(E'')} &\geq \frac{\dim V_m - \dim V'}{a_1(E'')} \\
&> \left(\dim V_m - h(m) \frac{a_1(F)}{a_1(h)} - \sum_i \varepsilon_i h_i(m) - \varepsilon \right) \frac{1}{a_1(E'')} \\
&= h(m) \frac{a_1(h) - a_1(F)}{a_1(h)} \frac{1}{a_1(E'')} - (\sum_i \varepsilon_i h_i(m) + \varepsilon) / a_1(E'') \\
&\geq h(m) \frac{a_1(h) - a_1(E')}{a_1(h)} \frac{1}{a_1(E'')} - \sum_i \varepsilon_i h_i(m) - \varepsilon \\
&= \frac{h(m)}{a_1(h)} - \sum_i \varepsilon_i h_i(m) - \varepsilon > 0.
\end{aligned}$$

There is a rational number λ_1 and an integer $m_0 \geq \lambda_1 - a_0(h)/a_1(h)$ which depend on $h(x)$, $\sum_i \varepsilon_i h_i(x)$ and ε such that

$$(7.32) \quad \frac{h(m)}{a_1(h)} - \sum_i \varepsilon_i h_i(m) - \varepsilon \geq \left(m + \frac{a_0(h)}{a_1(h)} - \lambda_1 \right)$$

for $m \geq m_0$.

By [S, Lem. 1.17], there is a purely 1-dimensional sheaf G of Hilbert polynomial $h(x)$ and a map $E \rightarrow G$ whose kernel is a coherent sheaf of dimension 0. Let $G \rightarrow G''$ be a quotient such that G'' is semi-stable. We set $E' := \ker(E \rightarrow G'')$ and $E'' := \text{im}(E \rightarrow G'')$. Since $h^0(X, E''(m)) \leq h^0(X, G''(m))$ and $a_1(E'') = a_1(G'')$,

$$\begin{aligned}
(7.33) \quad \frac{h^0(X, G''(m))}{a_1(G'')} &\geq \frac{h^0(X, E''(m))}{a_1(E'')} \\
&\geq \frac{h(m)}{a_1(h)} - \sum_i \varepsilon_i h_i(m) - \varepsilon > 0.
\end{aligned}$$

Since G'' is semi-stable, [S, Cor. 1.7] implies that

$$(7.34) \quad \frac{h^0(X, G''(m))}{a_1(G'')} \leq \begin{cases} 0, & c + m + \frac{\chi(G'')}{a_1(G'')} < 0 \\ c + m + \frac{\chi(G'')}{a_1(G'')}, & c + m + \frac{\chi(G'')}{a_1(G'')} \geq 0, \end{cases}$$

where c is a constant which only depends on $a_1(h)$. Since $h^0(X, G''(m)) > 0$, we get $c + m + \chi(G'')/a_1(G'') \geq 0$ and $\chi(G'')/a_1(G'') \geq a_0(h)/a_1(h) - \lambda_1 - c$, which means that G is of type $\lambda := a_0(h)/a_1(h) - \lambda_1 - c$. Replacing m , we may assume that for all type λ sheaves I of Hilbert polynomial $h(x)$, $I(m)$ is generated by global sections and $H^i(X, I(m)) = 0$, $i > 0$. In particular $h^0(X, G(m)) = h(m) = \dim V_m$. Assume that $H^0(X, E(m)) \rightarrow H^0(X, G(m))$ is not injective and let V' be the kernel. Then we get a contradiction from the inequality (7.29) and $\alpha_1 > 0$. Thus $H^0(X, E(m)) \rightarrow H^0(X, G(m))$ is injective, and hence it is isomorphic. Since $G(m)$ is generated by global sections, $E \rightarrow G$ must be surjective, which implies that it is isomorphic. Therefore E is of pure dimension d , of type λ and $V_m \rightarrow H^0(X, E(m))$ is an isomorphism. Thus we complete the proof of Claim 7.2.

We assume that $m \geq m(\lambda)$. Then $H^0(X, E(m)) \rightarrow H^0(X, E_i(m))$ is surjective. Thus $\beta_i : V \otimes R \rightarrow p_{T*}(\mathcal{E}_i(m))$ is surjective and define a morphism $T \rightarrow G_i$. Since $\beta_{i|U} = \alpha_{i|U}$ as elements of G_i , we get $\beta_i = \alpha_i$. Assume that there is a generalized parabolic quotient $E_* \rightarrow E''_*$ which destabilizes semi-stability. Since $E'_* := \ker(E_* \rightarrow E''_*)$ satisfies (b1, 2), we get that $V' = H^0(X, E'(m))$, $\alpha_i(V') = H^0(X, E'_i(m))$ and

$$(7.35) \quad \frac{\text{par-}\chi(E''_*(m))}{a_1(E'')} \geq \frac{\text{par-}\chi(E_*(m))}{a_1(h)} - \varepsilon.$$

Since ε_i are rational numbers, for a sufficiently small ε , we get

$$(7.36) \quad \frac{\text{par-}\chi(E''_*(m))}{a_1(E'')} \geq \frac{\text{par-}\chi(E_*(m))}{a_1(h)},$$

which is a contradiction. Therefore E is generalized parabolic semi-stable. Thus we get a lifting of $T \rightarrow \Gamma^{ss}$ and conclude that $\Gamma^{ss} \rightarrow (G(n) \times \prod_i G_i)^{ss}$ is proper. $\square$

By standard arguments, we see that $SL(V_m)s$, $s \in \Gamma^{ss}$ is a closed orbit if and only if the corresponding generalized parabolic semi-stable sheaf $(E, F(E), \alpha_*)$ is isomorphic to $\oplus_i (E_i, F(E_i), \alpha_*)$, where $(E_i, F(E_i), \alpha)$ are generalized parabolic stable sheaves.

Theorem 7.6. *There is a moduli scheme $\overline{M}_{X/\mathbb{C}}^{h_*, \alpha_*}$ parametrizing S -equivalence classes of generalized parabolic semi-stable sheaves $(E, F(E), \alpha_*)$ of Hilbert polynomials h_* .*

Let D be a Cartier divisor on X . Then

$$(7.37) \quad \{(E, F(E), \alpha_*) | E(-D) \rightarrow E \rightarrow E_l \text{ is a 0 map}\}$$

has a natural closed subscheme structure of $\overline{M}_{X/\mathbb{C}}^{h_*, \alpha_*}$. Assume that $h_l(m) = \chi(E(m)) - \chi(E(-D)(m))$. Then if $\dim(D \cap \text{Supp}(E)) = 0$, then $E(-D) \rightarrow E$ is injective and the image is $F_{l+1}(E)$. Thus $(E, F(E), \alpha_*)$ becomes a parabolic semi-stable sheaf. Since this condition is an open condition, we get $\overline{M}_{D/X/\mathbb{C}}^{h_*, \alpha_*}$ as a locally closed subscheme of $\overline{M}_{X/\mathbb{C}}^{h_*, \alpha_*}$.

Theorem 7.7. *Let D be a Cartier divisor on X and assume that $h_l(m) = \chi(E(m)) - \chi(E(-D)(m))$. Then there is a moduli scheme $\overline{M}_{D/X/\mathbb{C}}^{h_*, \alpha_*}$ parametrizing S -equivalence classes of parabolic semi-stable sheaves $(E, F(E), \alpha_*)$ of Hilbert polynomials h_* .*

Assume that X is a surface. Let γ be an element of $K(X)$ such that $\text{rk } \gamma = 0$ and $c_1(\gamma)$ is effective. We set $h(x) = (c_1(\gamma), \mathcal{O}_X(1))x + \chi(\gamma)$. Let D is a irreducible and reduced divisor such that $(D, \mathcal{O}_X(1)) > (c_1(\gamma), \mathcal{O}_X(1))$. Then $\overline{M}_{D/X/\mathbb{C}}^{h_*, \alpha_*}$ becomes compact. Therefore we get Theorem 7.1.

Remark 7.2. Although we assume that X is defined over a field k , we can easily generalize our construction to relative setting $X \rightarrow S$, where S is of finite type over k and $X \rightarrow S$ is projective.

Acknowledgement. I would like to thank M. Inaba for explaining the difficulty of constructing the moduli space of parabolic semi-stable sheaves.

REFERENCES

- [A-I-K] Altman, A., Iarrobino, A., Kleiman, S., *Irreducibility of the compactified Jacobian*, Real and complex singularities (Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976), pp. 1–12. Sijthoff and Noordhoff, Alphen aan den Rijn, 1977.
- [BBH1] Bartocci, C., Bruzzo, U., Hernández Ruipérez, D., *A Fourier-Mukai transform for stable bundles on K3 surfaces*, J. Reine Angew. Math. **486** (1997), 1–16
- [BBH2] Bartocci, C., Bruzzo, U., Hernández Ruipérez, D., *Existence of μ -stable vector bundles on K3 surfaces and the Fourier-Mukai transform*, Algebraic geometry (Catania, 1993/Barcelona, 1994), 245–257, Lecture Notes in Pure and Appl. Math., 200, Dekker, New York, 1998
- [Br1] Bridgeland, T., *Fourier-Mukai transforms for elliptic surfaces*, J. reine angew. Math. **498** (1998), 115–133
- [Br2] Bridgeland, T., *Equivalences of triangulated categories and Fourier-Mukai transforms*, Bull. London Math. Soc. **31** (1999), 25–34, math.AG/9809114
- [B-M] Bruzzo, U., Maciocia, A., *Hilbert schemes of points on some K3 surfaces and Gieseker stable bundles*, Math. Proc. Cambridge Philos. Soc. **120** (1996), 255–261
- [E-G] Ellingsrud, G., Göttsche, L., *Variation of moduli spaces and Donaldson invariants under change of polarization*, J. Reine Angew. Math. **467** (1995), 1–49
- [F-Q] Friedman, R., Qin, Z., *Flips of moduli spaces and transition formulas for Donaldson polynomial invariants of rational surfaces*, Comm. Anal. Geom. **3** (1995), 11–83
- [G] Gieseker, D., *On the moduli of vector bundles on an algebraic surface*, Ann. of Math. **106** (1977), 45–60
- [Gö] Göttsche, L., *Hilbert schemes of zero-dimensional subschemes of smooth varieties*, Lecture Notes in Mathematics, 1572. Springer-Verlag, Berlin, (1994)
- [Gö] Göttsche, L., *Theta functions and Hodge numbers of moduli spaces of sheaves on rational surfaces*, Comm. Math. Phys. **206** (1999), 105–136
- [G-H] Göttsche, L., Huybrechts, D., *Hodge numbers of moduli spaces of stable bundles on K3 surfaces*, Internat. J. Math. **7** (1996), 359–372
- [H-M] Hernández Ruipérez, D., Muñoz Porras, J. M., *Structure of the moduli space of stable sheaves on elliptic fibrations*, Preprint Universidad de Salamanca, (1998)
- [I] Inaba, M., *Moduli of parabolic stable sheaves on a projective scheme*, J. Math. Kyoto Univ. **40** (2000), 119–136
- [J-M] Jardim, M., Maciocia, M., *A Fourier-Mukai approach to spectral data for instantons*, math.AG/0006054
- [Li] Li, J., *Compactification of moduli of vector bundles over algebraic surfaces*, Collection of papers on geometry, analysis and mathematical physics, World Sci. Publishing, River Edge, NJ, (1997), 98–113
- [Ma1] Maruyama, M., *Moduli of stable sheaves II*, J. Math. Kyoto Univ. **18** (1978), 557–614
- [Ma2] Maruyama, M., *Moduli of algebraic vector bundles*, in preparation
- [Ma-Yk] Maruyama, M., Yokogawa, K., *Moduli of parabolic stable sheaves*, Math. Ann. **293** (1992), 77–99
- [M-W] Matsuki, K., Wentworth, R., *Mumford-Thaddeus principle on the moduli space of vector bundles on an algebraic surface*, Internat. J. Math. **8** (1997), 97–148
- [Mt] Matsushita, D., *Equidimensionality of Lagrangian fibrations on holomorphic symplectic manifolds*, Math. Res. Lett. **7** (2000), 389–391
- [MNVW] Minahan, J.A., Nemeschansky, D., Vafa, C., Warner, N.P., *E-Strings and $N = 4$ Topological Yang-Mills Theories*, Nucl. Phys. B **527** (1998), 581–623 hep-th/9802168
- [Mu1] Mukai, S., *Semi-homogeneous vector bundles on an Abelian variety*, J. Math. Kyoto Univ. **18** (1978), 239–272
- [Mu2] Mukai, S., *Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves*, Nagoya Math. J., **81** (1981), 153–175
- [Mu3] Mukai, S., *Symplectic structure of the moduli space of sheaves on an abelian or K3 surface*, Invent. math. **77** (1984), 101–116

- [Mu4] Mukai, S., *On the moduli space of bundles on K3 surfaces I*, Vector bundles on Algebraic Varieties, Oxford, 1987, 341–413
- [Mu5] Mukai, S., *Fourier functor and its application to the moduli of bundles on an Abelian variety*, Adv. Studies in Pure Math. **10** (1987), 515–550
- [Mu6] Mukai, S., *Moduli of vector bundles on K3 surfaces, and symplectic manifolds*, Sugaku Expositions, **1** (1988), 139–174
- [Mu7] Mukai, S., *Abelian variety and spin representation (in Japanese)*, Proceedings of symposium “Hodge theory and algebraic geometry (Sapporo, 1994)”, 110–135: English translation, Univ. of Warwick preprint, 1998
- [Mu8] Mukai, S., *Duality of polarized K3 surfaces*, Proceedings of Euroconference of Algebraic Geometry, 1996 to appear
- [MFK] Mumford, D., Fogarty, J., Kirwan, F., *Geometric invariant theory*, Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2) 34. Springer-Verlag, Berlin, 1994.
- [O1] O’Grady, K., *The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface*, J. Algebraic Geom. **6** (1997), 599–644
- [O2] O’Grady, K., *Desingularized moduli spaces of sheaves on a K3*, J. reine angew. Math. **512** (1999), 49–117, alg-geom/9708009, math.AG/9805099
- [O3] O’Grady, K., *A new six dimensional irreducible symplectic variety*, math.AG/0010187
- [Or] Orlov, D., *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. (NY), **84** (1997) 1361–1381
- [S] Simpson, C., *Moduli of representations of the fundamental group of a smooth projective variety I*, Publ. Math. I.H.E.S. **79** (1994), 47–129
- [V-W] Vafa, C., Witten, E., *A strong coupling test of S-duality*, Nucl. Phys. B **431** (1994), hep-th/9408074
- [Yk] Yokogawa, K., *Compactification of moduli of parabolic sheaves and moduli of parabolic Higgs sheaves*, J. Math. Kyoto Univ. **33** (1993), 451–504
- [Y1] Yoshioka, K., *Chamber structure of polarizations and the moduli of stable sheaves on a ruled surface*, Internat. J. Math. **7** (1996), 411–431
- [Y2] Yoshioka, K., *Numbers of $\mathbf{F}_q$ -rational points of the moduli of stable sheaves on elliptic surfaces*, Moduli of vector bundles (Sanda, 1994; Kyoto, 1994), 297–305, Lecture Notes in Pure and Appl. Math., 179, Dekker, New York, 1996.
- [Y3] Yoshioka, K., *Some notes on the moduli of stable sheaves on elliptic surfaces*, Nagoya Math. J. **154** (1999), 73–102
- [Y4] Yoshioka, K., *Some examples of Mukai’s reflections on K3 surfaces*, J. reine angew. Math. **515** (1999), 97–123
- [Y5] Yoshioka, K., *Euler characteristics of SU(2) instanton moduli spaces on rational elliptic surfaces*, Comm. Math. Phys. **205** (1999), 501–517
- [Y6] Yoshioka, K., *A note on moduli of stable sheaves on ruled surfaces*, Max-Planck-Institut für Mathematik Preprint series 1999 (104)
- [Y7] Yoshioka, K., *Irreducibility of moduli spaces of vector bundles on K3 surfaces*, math.AG/9907001
- [Y8] Yoshioka, K., *Moduli spaces of stable sheaves on abelian surfaces*, math.AG/0009001, Math. Ann. to appear

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KOBE UNIVERSITY, KOBE, 657, JAPAN
E-mail address: yoshioka@math.kobe-u.ac.jp